

LIMIT RELATIONS FOR THE COMPLEX ZEROS OF LAGUERRE AND q -LAGUERRE POLYNOMIALS

MARK V. DEFAZIO, DHARMA P. GUPTA, AND MARTIN E. MULDOON

ABSTRACT. For each $m (= 1, \dots, n)$ the n -th Laguerre polynomial $L_n^{(\alpha)}(x)$ has an m -fold zero at the origin when $\alpha = -m$. As the real variable $\alpha \rightarrow -m$, it has m simple complex zeros which approach 0 in a symmetric way. This symmetry leads to a finite value for the limit of the sum of the reciprocals of these zeros. There is a similar property for the zeros of the q -Laguerre polynomials and of the Jacobi polynomials and similar results hold for sums of other negative integer powers.

1. THE LAGUERRE POLYNOMIALS

The Laguerre polynomials are given by the explicit formula

$$(1) \quad L_n^{(\alpha)}(x) = \sum_{k=0}^n \binom{n+\alpha}{n-k} \frac{(-x)^k}{k!},$$

valid for all $x, \alpha \in \mathbf{C}$, showing them to be polynomials in both x and α .

We also use the representation [6, (5.3.3)],

$$(2) \quad L_n^{(\alpha)}(x) = \binom{n+\alpha}{n} {}_1F_1(-n, \alpha + 1; x) = \binom{n+\alpha}{n} \left[1 + \sum_{k=1}^n \frac{\binom{n}{k} (-x)^k}{(\alpha + 1)_k} \right],$$

with the usual notation for the confluent hypergeometric function. This is valid for $\alpha \neq -1, \dots, -n$, and with a limiting definition for $\alpha = -1, \dots, -n$.

The Laguerre polynomials satisfy the three term recurrence relation

$$(3) \quad xL_n^{(\alpha)}(x) = -(n+1)L_{n+1}^{(\alpha)}(x) + (\alpha + 2n + 1)L_n^{(\alpha)}(x) - (\alpha + n)L_{n-1}^{(\alpha)}(x),$$

with initial conditions $L_{-1}^{(\alpha)}(x) = 0$ and $L_0^{(\alpha)}(x) = 1$ for all complex α and x . When $\alpha > -1$, this recurrence relation is positive definite and the Laguerre polynomials are orthogonal with respect to the weight function $x^\alpha e^{-x}$ on $[0, +\infty)$. From this it follows that the zeros of $L_n^{(\alpha)}(x)$ are positive, real, simple, increasing functions of α

Date: January 3, 2007.

2000 Mathematics Subject Classification. 33C45, 33D45.

Key words and phrases. Laguerre polynomials, q -Laguerre polynomials, zeros, Jacobi polynomials.

This work was supported by grants from the Natural Sciences and Engineering Research Council, Canada. We thank a referee for pointing out some errors in an earlier version.

and they interlace with the zeros of $L_{n+1}^{(\alpha)}(x)$ [6]. When $\alpha \leq -1$ we no longer have orthogonality with respect to a positive weight function and the zeros can be non-real and non-simple. In the case $\alpha \leq -1$ the behaviour of the zeros of $L_n^{(\alpha)}(x)$ has been studied in detail in [1].

When $\alpha = -m$ ($1 \leq m \leq n$) the explicit formula (1) yields [6, (5.2.1)]

$$(4) \quad L_n^{(-m)}(x) = (-x)^m \frac{(n-m)!}{n!} L_{n-m}^{(m)}(x),$$

so $L_n^{(-m)}(x)$ has a zero of order m at $x = 0$, the remaining $n - m$ zeros being those of $L_{n-m}^{(m)}(x)$. The zero of order m arises from the coalescence of m simple zeros as α approaches $-m$. This follows by the method used in [6, §6.72] to prove some corresponding results for the Jacobi polynomials. (See [1] for more details of the Laguerre case.)

To see that m zeros approach the origin as $\alpha \rightarrow -m$, we let $\epsilon = \alpha + m$ and normalize so that the coefficient of x^m is 1; that is, we consider the following multiple of $L_n^{(\alpha)}(x)$:

$$(5) \quad q_n(x, \epsilon) = \sum_{k=0}^n c_k(\epsilon) x^k$$

where

$$(6) \quad c_k(\epsilon) = (-1)^{k-m} \frac{(n-m)! m! (\epsilon - m + k + 1)_{n-k}}{(n-k)! k! (\epsilon + 1)_{n-m}}.$$

For $0 \leq k \leq m-1$, $c_k(\epsilon)$ is a polynomial in ϵ which vanishes for $\epsilon = 0$; for $m-1 \leq k \leq n$, it is a rational function of ϵ which remains bounded as $\epsilon \rightarrow 0$; also $c_m(\epsilon) = 1$. Thus we may write

$$q_n(x, \epsilon) = x^m + \epsilon c'_0(0) + r(x, \epsilon),$$

where $r(x, \epsilon)$ remains to be examined more closely. If we choose an x_0 so that $|x_0| = 2|\epsilon c'_0(0)|^{1/m}$, we find that

$$r(x_0, \epsilon) = O(\epsilon^{1+1/m}), \quad \epsilon \rightarrow 0.$$

so comparing $r(x_0, \epsilon)$ and $x^m + \epsilon c'_0(0)$, we find, by Rouché's theorem, that $q_n(x, \epsilon)$ has exactly m zeros in the disk $|x| < |x_0|$. This shows that the m -fold zero of the right-hand side of (4) arises from the confluence of m zeros of $L_n^{(\alpha)}(x)$, as $\alpha \rightarrow -m$.

For $n \geq 2$, although the reciprocal of each zero becomes infinite, as $\alpha \rightarrow -m$, the sum of their reciprocals approaches a finite negative value. We state this as follows:

THEOREM 1. *Let $x_1(\alpha), \dots, x_m(\alpha)$ be the m ($2 \leq m \leq n$) zeros of $L_n^{(\alpha)}(x)$ in a neighbourhood of $x = 0$ for $\alpha \sim -m$. Then*

$$(7) \quad \lim_{\alpha \rightarrow -m} \sum_{k=1}^m \frac{1}{x_k(\alpha)} = \frac{m(m-2n-1)}{m^2-1}.$$

Proof: Let α be close to $-m$. Let us number the zeros of $L_n^{(\alpha)}(x)$ so that x_1, \dots, x_m are near zero and x_{m+1}, \dots, x_n are near the positive real zeros of $L_{n-m}^{(m)}(x)$. The explicit formula (2) yields

$$(8) \quad \sum_{k=1}^n \frac{1}{x_k(\alpha)} = \frac{n}{\alpha + 1}, \quad \alpha \neq -1, \dots, -n.$$

This is readily seen by considering the factorization

$$(9) \quad ({}^{n+\alpha}_n)^{-1} L_n^{(\alpha)}(x) = \left(1 - \frac{x}{x_1(\alpha)}\right) \cdots \left(1 - \frac{x}{x_n(\alpha)}\right) = 1 - x \sum_{k=1}^n \frac{1}{x_k(\alpha)} + \cdots,$$

and comparing with the expansion (2):

$$(10) \quad ({}^{n+\alpha}_n)^{-1} L_n^{(\alpha)}(x) = 1 - \frac{n}{\alpha + 1}x + \cdots.$$

Letting $\alpha \rightarrow -m$ gives

$$\begin{aligned} \frac{n}{1-m} &= \lim_{\alpha \rightarrow -m} \sum_{k=1}^n \frac{1}{x_k(\alpha)} \\ &= \lim_{\alpha \rightarrow -m} \sum_{k=1}^m \frac{1}{x_k(\alpha)} + \lim_{\alpha \rightarrow -m} \sum_{k=m+1}^n \frac{1}{x_k(\alpha)}, \end{aligned}$$

where the zeros in the first sum are the ones in the neighbourhood of 0. But the zeros in the second sum on the right approach those of $L_{n-m}^{(m)}(x)$ and hence, using (8), we get

$$(11) \quad \frac{n}{1-m} = \lim_{\alpha \rightarrow -m} \sum_{k=1}^m \frac{1}{x_k(\alpha)} + \frac{n-m}{1+m},$$

which gives (7).

2. q -LAGUERRE POLYNOMIALS

For $0 < q < 1$, the q -Laguerre polynomials may be defined by

$$(12) \quad L_n^{(\alpha)}(x, q) = \frac{(q^{\alpha+1}; q)_n}{(q; q)_n} \sum_{k=0}^n \frac{(q^{-n}; q)_k q^{k(k+1)/2+k(n+\alpha)}}{(q^{\alpha+1}; q)_k (q; q)_k} (1-q)^k x^k,$$

where we use the standard notation

$$(a; q)_n = \begin{cases} 1, & n = 0, \\ (1-a)(1-aq) \cdots (1-aq^{n-1}), & n = 1, 2, \dots \end{cases}$$

The definition (12) is that given by Moak [5] and in [2, p. 210]. The definition given in [4, p. 108] has x replaced by $(1-q)^{-1}x$.

Note that $L_n^{(\alpha)}(x, q) \rightarrow L_n^{(\alpha)}(x)$ as $q \rightarrow 1^-$ so if we use $x_k(\alpha; q)$ for the zeros of $L_n^{(\alpha)}(x, q)$ then $x_k(\alpha; q) \rightarrow x_k(\alpha)$, as $q \rightarrow 1^-$.

It is an easy matter to show that

$$(13) \quad L_n^{(-m)}(x, q) = (-x)^m (1-q)^m \frac{(q; q)_{n-m}}{(q; q)_n} L_{n-m}^{(m)}(x, q), \quad m = 1, 2, \dots, n.$$

We may proceed as in §1, to get

$$(14) \quad \sum_{k=1}^n \frac{1}{x_k(\alpha; q)} = \frac{q^{\alpha+1}(1-q^n)}{1-q^{\alpha+1}}, \quad \alpha \neq -1, \dots, -n,$$

which reduces to (8) as $q \rightarrow 1^-$, and

$$(15) \quad \lim_{\alpha \rightarrow -m} \sum_{k=1}^m \frac{1}{x_k(\alpha; q)} = \frac{q^{n-1}(1-q^{-m})(1+q-q^{m-n}-q^{-n})}{(1-q^{m-1})(1-q^{-m-1})}, \quad m = 2, \dots, n$$

which reduces to (7) as $q \rightarrow 1^-$.

3. SUMS OF POWERS OF THE ZEROS

Here we consider power sums

$$S_r = \sum_{k=1}^n x_k^{-r}, \quad r = 1, 2, \dots,$$

of the zeros of $L_n^{(\alpha)}(x)$. If we write

$$\binom{n+\alpha}{n}^{-1} L_n^{(\alpha)}(x) = \prod_{k=1}^n \left(1 - \frac{x}{x_k(\alpha)}\right) = 1 + \sum_{k=1}^n a_k x^k,$$

then the sums S_j are related to the coefficients a_j by [3, §3]:

$$S_1 = -a_1, \quad S_r = -ra_r - \sum_{i=1}^{r-1} a_i S_{r-i}, \quad r = 2, 3, \dots$$

In particular $S_2 = a_1^2 - 2a_2$, giving, in the Laguerre case,

$$S_2 = \frac{n^2 + n(\alpha + 1)}{(\alpha + 1)^2(\alpha + 2)}.$$

We may argue as in the proof of Theorem 1 that

$$\begin{aligned} \frac{n^2}{(1-m)^2} - \frac{n(n-1)}{(m-1)(m-2)} &= \lim_{\alpha \rightarrow -m} \sum_{k=1}^n \frac{1}{x_k(\alpha)^2} \\ &= \lim_{\alpha \rightarrow -m} \sum_{k=1}^m \frac{1}{x_k(\alpha)^2} + \lim_{\alpha \rightarrow -m} \sum_{k=m+1}^n \frac{1}{x_k(\alpha)^2}. \end{aligned}$$

As in §1, the zeros in the second sum on the right approach those of $L_{n-m}^{(m)}(x)$ and hence, with the notation of Theorem 1,

$$(16) \quad \frac{n^2}{(1-m)^2} - \frac{n(n-1)}{(m-1)(m-2)} = \lim_{\alpha \rightarrow -m} \sum_{k=1}^m \frac{1}{x_k(\alpha)^2} + \frac{(n-m)^2}{(1+m)^2} - \frac{(n-m)(n-m-1)}{(m+1)(m+2)},$$

This leads to

$$(17) \quad \lim_{\alpha \rightarrow -m} \sum_{k=1}^m \frac{1}{x_k(\alpha)^2} = -\frac{(n-m)(n+1)}{(m+1)^2(m+2)} - \frac{n(n-m+1)}{(m-1)^2(m-2)}, \quad 3 \leq m \leq n.$$

It makes sense that the above sum is meaningless for $m = 2$ since we do not expect the required cancellation in that case; $x_1(\alpha)^2$ and $x_2(\alpha)^2$ are positive numbers which both approach 0 as $\alpha \rightarrow -2^+$.

We may apply the same method to the q -Laguerre polynomials to get

$$(18) \quad \lim_{\alpha \rightarrow -m} \sum_{k=1}^m \frac{1}{x_k^2(\alpha; q)} = \frac{(1-q^n)^2}{(1-q^{m-1})^2} - \frac{2q(1-q^n)(1-q^{n-1})}{(1+q)(1-q^{m-1})(1-q^{m-2})} \\ - \frac{q^{2m+2}(1-q^{n-m})^2}{(1-q^{m+1})^2} + \frac{2q^{2m+4}(1-q^{n-m})(1-q^{n-m-1})}{(1+q)(1-q^{m+1})(1-q^{m+2})},$$

which reduces to (17) as $q \rightarrow 1^-$.

4. JACOBI POLYNOMIALS

The reason for the described behaviour of the zeros of Laguerre polynomials lies in their hypergeometric character as exhibited in (2). In fact our results may be extended to a general class of ${}_pF_q$ with a numerator parameter $-n$ and a denominator parameter $\alpha + 1$, even in some cases where the extra parameters depend on α . We confine attention to the case of Jacobi polynomials.

As in [6, (4.21.2)], we may define the Jacobi polynomial $P_n^{(\alpha, \beta)}(x)$, for general complex numbers α and β by

$$(19) \quad P_n^{(\alpha, \beta)}(x) = \frac{1}{n!} \sum_{k=0}^n \binom{n}{k} (n + \alpha + \beta + 1)_k (\alpha + k + 1)_{n-k} \left(\frac{x-1}{2} \right)^k,$$

or by

$$(20) \quad P_n^{(\alpha, \beta)}(x) = \binom{n+\alpha}{n} {}_2F_1 \left(-n, n + \alpha + \beta + 1; \alpha + 1; \frac{1-x}{2} \right),$$

with a limiting definition for $\alpha = -1, \dots, -n$, from which it follows [6, (4.22.2)] that, for $1 \leq m \leq n$,

$$(21) \quad \binom{n}{m} P_n^{(-m, \beta)}(x) = \binom{n+\beta}{m} \left(\frac{x-1}{2} \right)^m P_{n-m}^{(m, \beta)}(x).$$

From the explicit formula (20) or from the differential equation for the Jacobi polynomials, we have, for $\alpha \neq -1, \dots, -n$,

$$(22) \quad \sum_{k=1}^n \frac{1}{1 - x_k(\alpha, \beta)} = \frac{n(n + \alpha + \beta + 1)}{2(\alpha + 1)}.$$

From this we can get

THEOREM 2. *Let $x_1(\alpha, \beta), \dots, x_m(\alpha, \beta)$ be the m ($2 \leq m \leq n$) zeros of $P_n^{(\alpha, \beta)}$ in a neighbourhood of $x = 1$ for $\alpha \sim -m$. Then*

$$(23) \quad \lim_{\alpha \rightarrow -m} \sum_{k=1}^m \frac{1}{1 - x_k(\alpha, \beta)} = \frac{n(n - m + \beta + 1)}{2(1 - m)} - \frac{(n - m)(n + \beta + 1)}{2(m + 1)}.$$

Limits of sums of higher powers and q -extensions are obtainable as before.

REFERENCES

- [1] Mark V. DeFazio, *On the zeros of some quasi-definite orthogonal polynomials*, Ph.D. Dissertation, York University, 2001.
- [2] G. Gasper and M. Rahman, *Basic Hypergeometric Series*, 2nd ed., Cambridge University Press, 2004.
- [3] M. E. H. Ismail and M. E. Muldoon, Bounds for the small real and purely imaginary zeros of Bessel and related functions, *Meth. Appl. Anal.* **2** (1995), 1–21.
- [4] R. Koekoek and R. Swarttouw, *The Askey-scheme of hypergeometric orthogonal polynomials and its q -analogue*, Delft University of Technology, Department of Technical Mathematics and Informatics Report no. 98-17, 1998. <http://aw.twi.tudelft.nl/~koekoek/askey.html>
- [5] D. S. Moak, The q -analogue of the Laguerre polynomials. *J. Math. Anal. Appl.* **81** (1981), 20–47.
- [6] G. Szegő, *Orthogonal Polynomials*, Amer. Math. Soc. Colloq. Publ., vol. 23, 4th ed., Amer. Math. Soc., Providence, R.I., 1975.

ALGOMA UNIVERSITY COLLEGE, SAULT STE. MARIE, ONTARIO P6A 2G4, CANADA

DEPARTMENT OF MATHEMATICS AND STATISTICS, YORK UNIVERSITY, TORONTO, ONTARIO M3J 1P3, CANADA

DEPARTMENT OF MATHEMATICS AND STATISTICS, YORK UNIVERSITY, TORONTO, ONTARIO M3J 1P3, CANADA

E-mail address: muldoon@yorku.ca