

# Expansion of the Riemann $\Xi$ function in Meixner-Pollaczek polynomials \*

Alexey Kuznetsov †

Current version: February 2, 2007

## Abstract

In this article we study in detail the expansion of the Riemann  $\Xi$  function in Meixner-Pollaczek polynomials. We obtain explicit formulas, recurrence relation and asymptotic expansion for the coefficients and investigate the zeros of the partial sums.

---

\*To appear in the Canadian Mathematical Bulletin

†Research supported by the Natural Sciences and Engineering Research Council of Canada and MITACS Mathematics of Information Technology and Complex Systems, Canada. The first version of the manuscript was completed while the author was a Postdoctoral Fellow at the Department of Mathematics and Statistics, McMaster University. The author would like to thank an anonymous referee for many helpful comments.

# 1 Introduction

Riemann  $\Xi$  function is defined as

$$\Xi(t) = \frac{1}{2}s(s-1)\pi^{-\frac{s}{2}}\Gamma\left(\frac{s}{2}\right)\zeta(s), \quad s = \frac{1}{2} + it.$$

It is known that  $\Xi(t)$  is an even entire function, which is real for real  $t$ , and its zeros coincide with the nontrivial zeros of the Riemann zeta function  $\zeta\left(\frac{1}{2} + it\right)$  (see [11]).

In this article we study the expansion of the  $\Xi$ -function in the Meixner-Pollaczek polynomials. The expansions in polynomials are important due to the following reasons: first of all, the location of the zeros of partial sums  $S_n(t)$  can provide information about the zeros of the  $\Xi$ -function: for example, if the polynomials  $S_n(t)$  have only real zeros and converge uniformly on compact subsets of  $\mathbb{C}$  to  $\Xi(t)$ , then  $\Xi(t)$  itself has only real zeros. Second reason is that the zeros of the partial sums  $S_n(t)$  can be computed numerically quite easily (compared to the Dirichlet series or expansions in special functions). Third reason is that there are many results in the theory of multiplier sequences, biorthogonal polynomials and linear operators on polynomials that can potentially be applied to study the roots of the partial sums  $S_n(t)$ .

In order to be useful the expansion must be simple and natural: thus it is important to choose the right basis functions. Depending on the choice of the basis the coefficients of the expansion can be given by explicit formulas and be easily computable or they can be completely intractable. For example, the well known Stieltjes constants  $\gamma_n$  are defined (up to plus/minus sign) as the derivatives of  $\zeta(s) - \frac{1}{s-1}$  at  $s = 1$ . These numbers were studied extensively (see [13]), here we just note the following two formulas:

$$\gamma_n = \lim_{m \rightarrow \infty} \left[ \sum_{j=1}^m \frac{\ln(j)^n}{j} - \frac{\ln(m)^{n+1}}{n+1} \right] = \frac{\ln(2)^n}{n+1} \sum_{j \geq 1} \frac{(-1)^j}{j} B_{j+1} \left( \frac{\ln(j)}{\ln(2)} \right).$$

One can see that the above formulas are very complicated, it is hard to extract any information about  $\gamma_n$  from either of them. This indicates that  $(s-1)^n$  is not a very convenient basis for expansion.

The fact that power functions do not form a convenient basis for expansion is not surprising: zeta function is not a solution of any simple ODE. Close connection of zeta function with Mellin transform indicates that falling factorials  $\binom{s-1}{j}$  might be a better choice for basis functions. The following expansion of  $\zeta(s)$  can be obtained with the help of the Gauss-Kuzmin-Wirsing operator (see [3],[12])

$$\zeta(s) = \frac{s}{s-1} - \frac{1}{2} - s \sum_{n \geq 0} (-1)^n \binom{s-1}{n} a_n.$$

Coefficients  $a_n$  are given in terms of values of  $\zeta(s)$  at integers:

$$a_n = 1 - \frac{1}{2(n+1)} - \gamma + \sum_{k=1}^n (-1)^k \binom{n}{k} \left[ \frac{1}{k} - \frac{\zeta(k+1)}{k+1} \right].$$

Even though  $a_n$  are not given in closed form, they can be computed with high precision (at least for small  $n$ ).

The above expansions have one major drawback: their partial sums don't respect the functional equation. If one wants the zeros of the partial sums to be real, the partial sums  $S_n(t)$  themselves must be real for real  $t$  (up to a constant multiple), which means that they must satisfy the functional equation. It turns out that a natural basis for expansion of the  $\Xi$ -function is given by the Meixner-Pollaczek polynomials. In this article we show that the coefficients of this expansion have many important and useful properties: closed form expression, explicit formula for the exponential generating function, recurrence relation, integral representation and asymptotic expansion.

## 2 Expansion in Meixner-Pollaczek polynomials

In this article we consider only a special case of the Meixner-Pollaczek polynomials, defined as

$$p_n(t) = P_n^{(\frac{3}{4})} \left( \frac{t}{2}; \frac{\pi}{2} \right) = \frac{(2n+1)!!}{(2n)!!} i^n {}_2F_1 \left( -n, \frac{3}{4} + \frac{it}{2}; \frac{3}{2}; 2 \right).$$

For the definition and properties of the general Meixner-Pollaczek polynomials  $P_n^{(\lambda)}(t; \phi)$  see [8].

Polynomials  $p_n(t)$  arise as the Mellin transform of the odd Hermite orthogonal functions  $f_n(x) = e^{-\frac{x^2}{2}} H_{2n+1}(x)$ :

$$\int_0^{\infty} f_n(x) x^{s-1} dx = n! 2^{2n+\frac{s+1}{2}} \Gamma \left( \frac{1+s}{2} \right) i^n p_n(t), \quad s = \frac{1}{2} + it. \quad (1)$$

Since Hermite functions  $f_n(x)$  are invariant under Fourier transform, polynomials  $p_n(t)$  satisfy the "functional equation"  $p_n(-t) = (-1)^n p_n(t)$ . From the Parseval identity for the Mellin transform and (1) it follows that polynomials  $p_n(t)$  are orthogonal on  $\mathbb{R}$  with respect to the measure  $\mu(dt) = |\Gamma(\frac{3}{4} + \frac{it}{2})|^2 dt$ , which also implies that  $p_n(t)$  have only real zeros (see [1]). The three term recurrence relation for  $p_n(t)$  is

$$(n+1)p_{n+1}(t) - tp_n(t) + \left(n + \frac{1}{2}\right) p_{n-1}(t) = 0, \quad n \geq 0. \quad (2)$$

The next theorem shows that Meixner-Pollaczek polynomials  $p_n(t)$  form a natural basis for expansion of  $\Xi$ -function:

**Theorem 2.1.** *Define the sequence of real numbers  $\{\tilde{b}_n\}_{n \geq 0}$  as*

$$\tilde{b}_n = \frac{d^n}{dx^n} \left[ -\frac{1}{4} \sqrt{\frac{\pi}{x}} \sin \left( x + \frac{\pi}{8} \right) \tanh \left( \sqrt{2\pi x} \right) \right]_{x=0}. \quad (3)$$

Let the sequence  $\{b_n\}_{n \geq 0}$  be a linear transformation of  $\{\tilde{b}_n\}_{n \geq 0}$

$$b_n = 4n(n-1)\tilde{b}_{n-2} + (8n^2 + 12n + 7)\tilde{b}_n + (2n+3)(2n+5)\tilde{b}_{n+2}. \quad (4)$$

Then we have

$$\Xi(t)e^{-\frac{\pi t}{4}} = \sum_{n \geq 0} b_n p_n(t), \quad (5)$$

where the series converges uniformly on compact subsets of  $\mathbb{C}$ .

**Proof:** To prove this theorem we use the following result, derived in [9]: for all  $t \in \mathbb{C}$

$$\Xi(t)e^{-\frac{\pi t}{4}} = \frac{1}{2} \cos\left(\frac{\pi}{8}\right) - t \sin\left(\frac{\pi}{8}\right) + (1+4t^2) \sum_{n \geq 0} a_n p_n(t), \quad (6)$$

where the coefficients  $\{a_n\}_{n \geq 0}$  have the integral representation

$$a_n = \frac{(-1)^n}{(2n+1)!!} \int_0^\infty \frac{\sin\left(\frac{y^2}{2} + \frac{\pi}{8}\right)}{e^{2\sqrt{\pi}y} + 1} y^{2n+1} dy, \quad (7)$$

and the exponential generating function for  $\{a_n\}_{n \geq 0}$  is

$$\sum_{n \geq 0} \frac{a_n}{n!} x^n = \frac{1}{4} \sqrt{\frac{\pi}{x}} \left[ -\sin\left(x + \frac{\pi}{8}\right) \tanh\left(\sqrt{2\pi x}\right) + \operatorname{Im}\left(\Phi\left(e^{\frac{\pi i}{4}} \sqrt{x}\right) e^{ix + \frac{\pi i}{8}}\right) \right]. \quad (8)$$

Here  $\Phi(x) = \frac{2}{\sqrt{\pi}} \int_0^x e^{-t^2} dt$  is the probability integral function (see [4]).

In order to obtain (5) all that is left to do is to express function  $(1+4t^2) \sum_{n \geq 0} a_n p_n(t)$  in (6) as expansion in  $p_n(t)$ : this will be done using the three term recurrence relation for the Meixner-Pollaczek polynomials  $p_n(t)$ .

Applying the recurrence relation (2) twice we find

$$(1+4t^2)p_n(t) = 4(n+1)(n+2)p_{n+2}(t) + (8n^2 + 12n + 7)p_n(t) + (2n-1)(2n+1)p_{n-2}(t). \quad (9)$$

Now, using (8) and (3) we find that for  $n \geq 0$

$$a_n - \tilde{b}_n = \frac{d^n}{dx^n} \left[ \frac{1}{4} \sqrt{\frac{\pi}{x}} \operatorname{Im}\left(\Phi\left(e^{\frac{\pi i}{4}} \sqrt{x}\right) e^{ix + \frac{\pi i}{8}}\right) \right],$$

thus

$$a_n = \tilde{b}_n + \frac{1}{2} \frac{(2n)!!}{(2n+1)!!} \cos\left(\frac{\pi}{2}n - \frac{\pi}{8}\right), \quad n \geq 0. \quad (10)$$

Next we use (9), (4) and (10) to transform the infinite sum in equation (6)

$$\begin{aligned} (1 + 4t^2) \sum_{n \geq 0} a_n p_n(t) &= \sum_{n \geq 0} p_n(t) [4n(n-1)a_{n-2} + (8n^2 + 12n + 7)a_n + (2n+3)(2n+5)a_{n+2}] = \\ &= -\frac{1}{2} \cos\left(\frac{\pi}{8}\right) + t \sin\left(\frac{\pi}{8}\right) + \sum_{n \geq 0} b_n p_n(t). \end{aligned}$$

Combining the above equation with (6) we obtain the required expansion (5). □

**Remark 1:** Expansion (6) was derived in [9] from the integral representation for  $\Xi(t)e^{-\frac{\pi t}{4}}$  by expanding the integral kernel (confluent hypergeometric function) in a series of Meixner-Pollaczek polynomials. A more intuitive way to derive (6) is to note that the fractional Fourier sine transform  $F(y) = [\mathcal{F}_s^{\frac{1}{2}} f](y)$  of the function  $f(x) = (\exp(\sqrt{2\pi}x) - 1)^{-1}$  can be computed in closed form (see [4],[9]), and  $F(y)$  can be used to obtain the explicit expression for the coefficients  $c_n$  of the following expansion

$$f\left(e^{\frac{\pi i}{4}} x\right) = \sum_{n \geq 0} c_n e^{-\frac{x^2}{2}} H_{2n+1}(x).$$

To obtain (6) we apply the Mellin transform to both sides of the above equation and use (1).

**Remark 2:** Note that for  $n \geq 2$  we have

$$b_n = 4n(n-1)a_{n-2} + (8n^2 + 12n + 7)a_n + (2n+3)(2n+5)a_{n+2} \quad (11)$$

and that linear transformation (4) eliminates the term  $\frac{1}{2} \frac{(2n)!!}{(2n+1)!!} \cos\left(\frac{\pi}{2}n - \frac{\pi}{8}\right)$  in (10).

**Remark 3:** The asymptotic formula for the Meixner-Pollaczek polynomials  $p_n(t) = P_n^{\left(\frac{3}{4}\right)}\left(\frac{t}{2}; \frac{\pi}{2}\right)$  can be derived from the integral representation (see [9]):

$$P_n^{(\lambda)}\left(t; \frac{\pi}{2}\right) \sim 2^{1-2\lambda} (-i)^n \left[ \frac{(2n)^{\lambda+it-1}}{\Gamma(\lambda+it)} + (-1)^n \frac{(2n)^{\lambda-it-1}}{\Gamma(\lambda-it)} \right], \quad n \rightarrow \infty. \quad (12)$$

We see that expansion in Meixner-Pollaczek polynomials converge in some strip  $\text{Im}(t) < c$ , and since  $\Xi(t)e^{-\frac{\pi t}{4}}$  is an entire function this expansion should converge in the entire complex plane. Another proof can be given with the help of the asymptotic formula for  $a_n$  derived in section 3.

Theorem 2.1 allows us to obtain an interesting expression for the nontrivial zeros of the Riemann zeta function as the limit of eigenvalues of some  $n \times n$  real (nonsymmetric) matrices

$\mathbf{B}_n$ , defined as

$$\mathbf{B}_n = \begin{pmatrix} 0 & 1 & 0 & \dots & 0 & 0 & 0 & 0 \\ \frac{3}{2} & 0 & 2 & \dots & 0 & 0 & 0 & 0 \\ 0 & \frac{5}{2} & 0 & \dots & 0 & 0 & 0 & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & \dots & 0 & n-3 & 0 & 0 \\ 0 & 0 & 0 & \dots & n-\frac{5}{2} & 0 & n-2 & 0 \\ 0 & 0 & 0 & \dots & 0 & n-\frac{3}{2} & 0 & n-1 \\ -n\frac{b_0}{b_n} & -n\frac{b_1}{b_n} & -n\frac{b_2}{b_n} & \dots & -n\frac{b_{n-4}}{b_n} & -n\frac{b_{n-3}}{b_n} & n-\frac{1}{2} & -n\frac{b_{n-2}}{b_n} & -n\frac{b_{n-1}}{b_n} \end{pmatrix} \quad (13)$$

Let  $\sigma(\zeta)$  denote the nontrivial zeros of the Riemann zeta function  $\zeta\left(\frac{1}{2} + it\right)$ :

$$\sigma(\zeta) = \{\gamma \in \mathbb{C} : \zeta\left(\frac{1}{2} + i\gamma\right) = 0, \text{Im}[\gamma] < \frac{1}{2}\}.$$

**Lemma 2.2.** *As  $n \rightarrow \infty$  the spectrum of matrices  $\mathbf{B}_n$  converges to the set of nontrivial zeros of the Riemann zeta function  $\zeta\left(\frac{1}{2} + it\right)$ :*

$$\sigma(\mathbf{B}_n) \longrightarrow \sigma(\zeta).$$

*The convergence is understood in the following sense:  $\gamma \in \sigma(\zeta)$  if and only if  $\gamma \neq \infty$  is a limit of some sequence  $\{\gamma_n\}_{n \geq 0}$ , such that  $\gamma_n \in \sigma(\mathbf{B}_n)$ .*

**Proof:** The proof is just an application of the Theorem 2.3 in [2] to the partial sums  $S_n(t) = \sum_{k=0}^n b_k p_k(t)$ . By Theorem 2.1 the partial sums  $S_n(t)$  converge to  $\Xi(t)e^{-\frac{\pi t}{4}}$  uniformly on compact subsets of  $\mathbb{C}$ , thus the zeros of  $S_n(t)$  converge to the set of zeros of  $\Xi(t)e^{-\frac{\pi t}{4}}$ , which coincides with  $\sigma(\zeta)$ . Now, to finish the proof we only need to use the Theorem 2.3 in [2], which allows us to express the roots of the polynomials  $S_n(t)$  as the eigenvalues of the matrix  $\mathbf{B}_n$  defined by (13). Here is the main idea: one can check using the recurrence relation (2) that

$$\mathbf{B}_n \mathbf{f}_n(t) = t \mathbf{f}_n(t) - \frac{n}{b_n} S_n(t) \mathbf{e}_n,$$

where  $\mathbf{f}_n(t) = [p_0(t), p_1(t), \dots, p_{n-1}(t)]^T$  and  $\mathbf{e}_n = [0, 0, \dots, 0, 1]^T$ . Thus  $S_n(\gamma) = 0$  if and only if  $\gamma$  is an eigenvalue of  $\mathbf{B}_n$  (and in this case  $\mathbf{f}_n(\gamma)$  is the right eigenvector). □

### 3 Computation of the coefficients

From equation (3) we can derive an explicit formula for the coefficients  $\tilde{b}_n$ :

$$\tilde{b}_n = -\frac{n!}{\sqrt{2}} \sum_{k=0}^n \frac{4^k (4^{k+1} - 1) B_{2k+2}}{(n-k)! (2k+2)!} \sin\left(\frac{\pi}{2}(n-k) + \frac{\pi}{8}\right) (2\pi)^{k+1}, \quad (14)$$

where  $\{B_n\}_{n \geq 0} = \{1, -\frac{1}{2}, \frac{1}{6}, 0, -\frac{1}{30}, 0, \frac{1}{42}, 0, \dots\}$  are the Bernoulli numbers. Coefficients  $\tilde{b}_n$  also satisfy the recurrence relation

$$\sum_{k=0}^n \frac{\pi^{n-k}}{(2n-2k-1)!!} \binom{n}{k} \tilde{b}_k = \frac{\sqrt{\pi} 2^{-n-2}}{(2n+1)!!} \operatorname{Im} \left[ e^{\frac{\pi i}{8}} i^n H_{2n+1} \left( \sqrt{\frac{\pi}{2}} e^{-\frac{\pi i}{4}} \right) \right]. \quad (15)$$

This relation can be obtained from the following equation for the generating function of  $\tilde{b}_n$ :

$$\cosh(\sqrt{2\pi x}) \sum_{n \geq 0} \frac{\tilde{b}_n}{n!} x^n = -\frac{1}{4} \sqrt{\frac{\pi}{x}} \sin\left(x + \frac{\pi}{8}\right) \sinh(\sqrt{2\pi x}),$$

which is derived from (3).

Equations (14) and (15) can be used to compute coefficients  $b_n$  for small  $n$ , however they are not efficient when  $n$  is large: the precision is lost in (14) because Bernoulli numbers and factorials grow very rapidly while recurrence relation (15) is unstable. Fortunately for large  $n$  one can use equation (11) combined with the integral representation (7) for  $a_n$  to compute the coefficients  $b_n$ . The idea is to use the saddle point method: first we rewrite (7) as

$$a_n = \operatorname{Im} \left[ \frac{(-1)^n e^{\frac{\pi i}{8}}}{(2n+1)!!} \int_0^\infty \frac{e^{i\frac{y^2}{2}}}{e^{2\sqrt{\pi}y} + 1} y^{2n+1} dy \right].$$

Note that the integral in the above equation grows as  $2^n n!$  as  $n \rightarrow \infty$ , but by shifting the contour of integration to  $e^{\frac{\pi i}{4}} \mathbb{R}_+$ , changing the variable of integration  $y \mapsto \sqrt{2n\pi} e^{\frac{\pi i}{4}} v$  and rearranging the terms we isolate the dominant term  $\exp(-2\sqrt{\pi n})$  and obtain more manageable expression:

$$\begin{aligned} a_n &= (-1)^n \frac{\sqrt{\pi}}{2} e^{-2\sqrt{\pi n}} e^{(n+1)\log(n) - n - \log(\Gamma(n+\frac{3}{2}))} \times \\ &\times \operatorname{Im} \left[ e^{\frac{\pi i}{8}} i^{n+1} e^{-2\sqrt{\pi n} i} \int_0^\infty \frac{e^{n(1-v+\log(v)) + \theta\sqrt{n}(\sqrt{v}-1)}}{1 + e^{\theta\sqrt{n}\sqrt{v}}} dv \right], \end{aligned} \quad (16)$$

where  $\theta = -2\sqrt{\pi}(1+i)$ . The integral in (16) has a saddle point essentially at  $v = 1$ , thus we introduce a new variable  $v = 1 + n^{-\frac{1}{2}}u$  and obtain:

$$a_n = (-1)^n \frac{\sqrt{\pi}}{2} e^{-2\sqrt{\pi n}} e^{(n+\frac{1}{2})\log(n) - n - \log(\Gamma(n+\frac{3}{2}))} \operatorname{Im} \left[ e^{\frac{\pi i}{8}} i^{n+1} e^{-2\sqrt{\pi n} i} I_n \right], \quad (17)$$

where

$$I_n = \int_{-\sqrt{n}}^\infty \frac{\exp \left[ n \left( \log \left( 1 + un^{-\frac{1}{2}} \right) - un^{-\frac{1}{2}} \right) + \theta n^{\frac{1}{2}} \left( \sqrt{1 + un^{-\frac{1}{2}}} - 1 \right) \right]}{1 + \exp \left[ \theta n^{\frac{1}{2}} \sqrt{1 + un^{-\frac{1}{2}}} \right]} du. \quad (18)$$

Now we can compute with high precision  $I_n$  (and thus  $a_n$  and  $b_n$  through formulas (17) and (11)) using numerical integration to evaluate the integral in (18). We even go one step further and derive an asymptotic formula for  $I_n$ : first we expand the integrand around  $u = 0$ :

$$\begin{aligned} & n \left( \log \left( 1 + un^{-\frac{1}{2}} \right) - un^{-\frac{1}{2}} \right) + \theta n^{\frac{1}{2}} \left( \sqrt{1 + un^{-\frac{1}{2}}} - 1 \right) = \\ & = -\frac{u^2}{2} + \frac{\theta u}{2} + n^{-\frac{1}{2}} \left( -\frac{\theta u^2}{8} + \frac{u^3}{3} \right) + n^{-1} \left( \frac{\theta u^3}{16} - \frac{u^4}{4} \right) + O \left( n^{-\frac{3}{2}} \right), \end{aligned}$$

and  $\left( 1 + \exp \left[ \theta n^{\frac{1}{2}} \sqrt{1 + un^{-\frac{1}{2}}} \right] \right)^{-1} = 1 + O \left( e^{-2\sqrt{n\pi}} \right)$ . Collecting all the terms we obtain

$$\begin{aligned} I_n &= \int_{-\infty}^{\infty} e^{-\frac{u^2}{2} + \frac{\theta u}{2}} \left[ 1 + \frac{1}{\sqrt{n}} \left( -\frac{\theta u^2}{8} + \frac{u^3}{3} \right) \right. \\ &\quad \left. + \frac{1}{n} \left( \frac{u^6}{18} - \frac{\theta u^5}{24} + \left( \frac{\theta^2}{128} - \frac{1}{4} \right) u^4 + \frac{\theta u^3}{16} \right) + O \left( n^{-\frac{3}{2}} \right) \right] du = \\ &= -\sqrt{2\pi} \left[ 1 + \sqrt{\frac{\pi}{n}} \left( \frac{(1-i)\pi}{6} - \frac{3(1+i)}{4} \right) + \frac{1}{n} \left( -\frac{\pi^3 i}{36} - \frac{\pi^2}{4} + \frac{7\pi i}{16} + \frac{1}{12} \right) + O \left( n^{-\frac{3}{2}} \right) \right] \end{aligned}$$

## 4 Zeros of expansions in orthogonal polynomials

An interesting approach to study zeros of expansions in orthogonal polynomials was developed in [5], [6], [7] and is based on the theory of biorthogonal polynomials. The following statement is a particular case of proposition 6 in [7]:

**Proposition 4.1.** *If a polynomial  $\hat{S}_n(x) = \sum_{k=0}^n \frac{(2k+1)!!}{(2k)!!} b_k x^k$  has only real zeros, then polynomial*

$$S_n(t) = \sum_{k=0}^n b_k p_k(t) \text{ also has only real zeros.}$$

Using the above proposition one can easily prove the following statement:

**Proposition 4.2.** *The entire function  $\tilde{\Xi}(t) = \sum_{n \geq 0} \frac{1}{(2n+1)!!} \tilde{b}_n p_n(t)$  has only real zeros.*

Note that  $\tilde{\Xi}(t)$  is “almost”  $\Xi(t)e^{-\frac{\pi t}{4}}$  (see equation (5)), except for the factor  $(2n+1)!!$  and the linear transformation (4). To prove proposition 4.2 we use (3) to show that the exponential generating function for  $\tilde{b}_n$

$$\sum_{n \geq 0} \frac{\tilde{b}_n}{n!} x^n = -\frac{1}{4} \sqrt{\frac{\pi}{x}} \sin \left( x + \frac{\pi}{8} \right) \tanh \left( \sqrt{2\pi x} \right)$$

is an entire function with only real zeros and we can approximate it uniformly by polynomials with only real zeros (for example, we can expand  $\sin \left( x + \frac{\pi}{8} \right)$ ,  $\sinh \left( \sqrt{2\pi x} \right)$  and  $\cosh \left( \sqrt{2\pi x} \right)$  as



infinite products of linear factors  $x - x_n$  and truncate the products). Applying proposition 4.1 to these polynomials completes the proof.

Thus if numerical computations would indicate that the partial sums  $S_n(t) = \sum_{k=0}^n b_k p_k(t)$  have only real zeros, proposition 4.1 and explicit formulas for  $b_n$  could provide a way to prove this; note, however, that proposition 4.1 gives only sufficient condition. Unfortunately, the polynomial  $S_3(t)$  already has two complex roots and the numerical computations indicate that the number of complex roots of  $S_n(t)$  increases as  $n$  increases.

However,  $S_n(t) = \sum_{k=0}^n b_k p_k(t)$  is not the only possible way to truncate the series (5). As we will see the situation can be improved if we use a *multiplier sequence*. A multiplier sequence  $\{m_k\}_{k \geq 0}$  is characterized by the following property: if a polynomial  $Q(x) = q_0 + q_1 x + \dots + q_n x^n$  has only real zeros, then  $T[Q](x) = m_0 q_0 + m_1 q_1 x + \dots + m_n q_n x^n$  also has only real zeros. A classical example of a multiplier sequence is given by the *Jensen sequence*

$$m_k^{(n)} = \frac{n!}{(n-k)!n^k} = \left(1 - \frac{1}{n}\right) \left(1 - \frac{2}{n}\right) \dots \left(1 - \frac{n-k+1}{n}\right),$$

note that  $m_k^{(n)} \rightarrow 1$  as  $n \rightarrow \infty$  and that  $m_k^{(n)} = 0$  if  $k > n$ . This sequence provides a “right” way to truncate Taylor series: a theorem by Jensen (see [10]) says that (under some additional conditions) an entire function  $f(x) = \sum_{k \geq 0} f_k x^k$  has only real zeros if and only if all polynomials  $T_n[f](x) = \sum_{k \geq 0} m_k^{(n)} f_k x^k$  have only real zeros.

Let’s consider for example the exponential function  $e^x$ : if we simply truncate the Taylor expansion  $Q_n(x) = \sum_{k=0}^n \frac{x^k}{k!}$ , then  $Q_n(x)$  converge to  $e^x$  uniformly on compact subsets of  $\mathbb{C}$  but it can be proven that all the zeros of  $Q_n(x)$  are complex (except for one real negative root when  $n$  is odd). However if we truncate the Taylor series in a different way, by using the Jensen multiplier sequence  $m_k^{(n)}$ , we obtain  $\tilde{Q}_n(x) = T_n[e^x](x) = \sum_{k=0}^n m_k^{(n)} \frac{x^k}{k!} = \left(1 + \frac{x}{n}\right)^n$ , and  $\tilde{Q}_n(x)$  has only real zeros. Note that since  $m_k^{(n)} \rightarrow 1$  as  $n \rightarrow \infty$ , polynomials  $Q_n(x)$  and  $\tilde{Q}_n(x)$  both converge to the same function, but  $\tilde{Q}_n(x)$  has more predictable behavior of its zeros: roots of  $\tilde{Q}_n(x)$  are real and converge to the root of  $e^x$  at  $x = -\infty$ .

If we naively apply the multiplier sequence  $m_k^{(n)}$  to the expansion of  $\Xi(t)e^{-\frac{\pi t}{4}}$  in Meixner-Pollaczek polynomials:

$$\tilde{S}_n(t) = T_n \left[ \Xi(t)e^{-\frac{\pi t}{4}} \right] (t) = \sum_{k=0}^n m_k^{(n)} b_k p_k(t),$$

then  $\tilde{S}_n(t)$  also converges to  $\Xi(t)e^{-\frac{\pi t}{4}}$  uniformly on compact subsets (probably at a much slower rate), but the roots of  $\tilde{S}_n(t)$  seem to behave more predictably: numerical computations show that all the roots of  $\tilde{S}_n(t)$  are real for  $n \leq 5$ , and  $\tilde{S}_n(t)$  has just two complex zeros for  $6 \leq n \leq 20$ . Of course the sequence  $m_k^{(n)}$  was designed to work with Taylor expansions; there

is no obvious reason why it should improve the behavior of roots for expansions in Meixner-Pollaczek polynomials. This seems to be an interesting question for the future work: to study the multiplier sequences and more general linear operators on expansions of entire functions in Meixner-Pollaczek polynomials and to investigate the behavior of zeros of the partial sums under such transformations.

## References

- [1] D. Bump, K.-K. Choi, P. Kurlberg and J. Vaaler *A local Riemann hypothesis, I.* Math. Zeitschrift. 233(2000), no. 1, 1-19.
- [2] D. Day and L. Romero, *Roots of polynomials expressed in terms of orthogonal polynomials.* SIAM J. Numer. Anal. 43, 5 (2005), 1969-1987.
- [3] A.Yu. Eremin, I.E. Kaporin and M.K. Kerimov, *The calculation of the Riemann zeta-function in the complex domain.* USSR Comput. Math. and Math. Phys. 25 (1985), no. 2, 111-119.
- [4] I.S. Gradshteyn and I.M. Ryzhik, *Tables of integrals, series and products.* 6-th edition, (2000).
- [5] A. Iserles, and S.P. Norsett. Zeros of transformed polynomials. SIAM J. Math. Anal. Vol. 21, No. 2 (1990), 483-509.
- [6] A. Iserles, and S.P. Norsett. On the theory of bi-orthogonal polynomials. Trans. Amer. Math. Soc 306 (1988), 455-474.
- [7] A. Iserles, and E.B. Saff. Zeros of expansions in orthogonal polynomials. Math. Proc. Camb. Phil. Soc. 105 (1989), 559-573.
- [8] R. Koekoek and R.F. Swarttouw, *The Askey-scheme of orthogonal polynomials and its q-analog.* Delft University of Technology, Faculty of Information Technology and Systems, Dep-t of Technical Mathematics and Informatics, Report no. 98-17, (1998).
- [9] A. Kuznetsov, *Integral representations for the Dirichlet L-functions and their expansions in Meixner-Pollaczek polynomials and Pochhammer functions.* Preprint, available online at <http://people.unb.ca/~akuznets>.
- [10] B.Ja. Levin, *Distribution of zeros of entire functions.* Transl. Math. Mono. vol. 5, Amer. Math. Soc., Providence, RI, (1964); revised ed (1980).
- [11] E.C. Titchmarsh, *The theory of the Riemann zeta-function.* Oxford University Press, 2nd ed. (1986).

- [12] L. Vepstas, *A series representation of the Riemann zeta derived from the Gauss-Kuzmin-Wirsing operator*. preprint, (2005).
- [13] E.W. Weisstein, *CRC concise encyclopedia of mathematics*. CRC Press, 2nd ed. (2002).

Department of Mathematical Sciences  
University of New Brunswick  
Saint John  
NB, E2L 4L5  
Canada  
e-mail: akuznets@unbsj.ca