

Analytic Hausdorff gaps

Ilijas Farah

In [6], Hausdorff constructed two families \mathcal{A} and \mathcal{B} of sets of integers such that

- (a) $A \cap B$ is finite for all $A \in \mathcal{A}$ and all $B \in \mathcal{B}$,
- (b) for every $C \subseteq \mathbb{N}$ either $A \setminus C$ is infinite for some $A \in \mathcal{A}$ or $B \cap C$ is infinite for some $B \in \mathcal{B}$, and
- (c) both \mathcal{A} and \mathcal{B} have order-type equal to ω_1 with respect to the inclusion modulo finite.

Families satisfying (a) are *orthogonal*, those satisfying both (a) and (b) form a *gap* in $\mathcal{P}(\mathbb{N})/\text{Fin}$ (or, they are *not separated* over Fin , the ideal of finite sets), and if they furthermore satisfy (c) they form an (ω_1, ω_1) -*gap*. If (a) and (b) hold and both \mathcal{A} and \mathcal{B} are linearly ordered by the inclusion modulo finite, a gap is *linear*. Before Hausdorff, du Bois-Reymond and Hadamard proved that there are no (ω, ω) -gaps in $\mathcal{P}(\mathbb{N})/\text{Fin}$ (see e.g., [11]). If (c) is weakened to (recall that a poset is *ω -directed* if every countable subset has an upper bound)

- (c') Both \mathcal{A} and \mathcal{B} are ω -directed under the inclusion modulo finite,

then the gap is said to be *Hausdorff* (note that a Hausdorff gap need not be linear).

We will identify subsets of \mathbb{N} with elements of the Cantor cube $2^{\mathbb{N}}$, and import the compact metric topology, as well as Borel structure, of the latter structure to $\mathcal{P}(\mathbb{N})$. Via this identification, the existence of an (ω_1, ω_1) -gap in $\mathcal{P}(\mathbb{N})/\text{Fin}$ has some immediate consequences to the classical descriptive set theory—e.g., it gives an example of an uncountable universal measure zero set (see [11, §4]) and a partition of the Cantor set into \aleph_1 many pairwise disjoint G_δ sets. More recently, Hausdorff gaps in $\mathcal{P}(\mathbb{N})/\text{Fin}$ have played a prominent role in a variety of subjects, from automatic continuity in Banach algebras ([3]) to the study of homomorphisms of $\mathcal{P}(\mathbb{N})/\text{Fin}$ (see [12, §IV], also [5, Chapter 3]).

In Hausdorff's time the fact that both sides of the gap have length equal to ω_1 probably could have been considered as a support for the Continuum Hypothesis. After the profusion of the method of forcing this specific property of ω_1 is better explained. In ([7]) Kunen has proved that, assuming a small amount of reflection at the level of $\mathcal{P}(\mathbb{N})$, (ω_1, ω_1) gaps are the only linear Hausdorff gaps in $\mathcal{P}(\mathbb{N})/\text{Fin}$. Since this reflection principle (later incorporated into Todorcevic's Open Coloring Axiom, see [16, §8]) is compatible with Martin's Axiom, ω_1 is the only standard

The author acknowledges support received from the National Science Foundation (USA) via grant DMS-0070798 and from the PSC-CUNY grant #62785-00-31.

cardinal invariant \mathfrak{d} of the continuum such that $\mathcal{P}(\mathbb{N})/\text{Fin}$ always has $(\mathfrak{d}, \mathfrak{d})$ - or (\mathfrak{d}, ω_1) -gaps. An important consequence of Kunen’s proof was that every Hausdorff gap can be made *indestructible* under any further \aleph_1 -preserving forcing; this fact has played a key role in Woodin’s consistency proof of ‘all homomorphisms of Banach algebras are automatically continuous’ (see [3]).

Inspired by these results, Todorćević proved a separation principle implying that there are no analytic Hausdorff gaps in $\mathcal{P}(\mathbb{N})/\text{Fin}$ ([17, Corollary 1]; see also [5, §5.5]). (Recall that a subset of a compact metric space is *analytic* if it is a continuous image of the irrationals.) His proof shows that in Solovay’s model ([15]) there are no Hausdorff gaps in $\mathcal{P}(\mathbb{N})/\text{Fin}$ at all, thus indicating that some form of Axiom of Choice is necessary to construct a Hausdorff gap in $\mathcal{P}(\mathbb{N})/\text{Fin}$. This may be the right place to point out to a rather optimal construction of a Hausdorff gap—also due to Todorćević—given in [2, pages 96–98]. The gap is dened by a simple formula whose parameters are an ω_1 -chain in $\mathcal{P}(\mathbb{N})/\text{Fin}$ and a ladder system on ω_1 .

If \mathcal{I} is an ideal on \mathbb{N} other than Fin , we say that two families \mathcal{A}, \mathcal{B} are \mathcal{I} -*orthogonal* (or *orthogonal over \mathcal{I}*) if $A \cap B \in \mathcal{I}$ for all $A \in \mathcal{A}$ and all $B \in \mathcal{B}$. The notions ‘separated over \mathcal{I} ’ and ‘(Hausdorff) gap in $\mathcal{P}(\mathbb{N})/\mathcal{I}$ ’ are dened in a natural way. The question of computing the gap-spectra of quotients other than $\mathcal{P}(\mathbb{N})/\text{Fin}$ was rst raised by W. Just (see [11]), and some partial answers were obtained by K. Mazur in [9].

The interest in the structure of analytic ideals and their quotients was intensified by [17] and its sequels [19, 18]. In [19] it was proved that every embedding of $\mathcal{P}(\mathbb{N})/\text{Fin}$ into a quotient over an analytic ideal that has a continuous (equivalently, Baire-measurable) lifting (see [5, §1.5]) preserves all Hausdorff gaps. Since by a result of Mathias ([8]) there is always such an embedding, this implies that the Hausdorff-gap spectrum of any analytic quotient $\mathcal{P}(\mathbb{N})/\mathcal{I}$ is at least as rich as the Hausdorff-gap spectrum of $\mathcal{P}(\mathbb{N})/\text{Fin}$. In particular, this implies the existence of (ω_1, ω_1) -gaps in every analytic quotient, answering the original Just’s question. (For more on preservation of gaps see [19] and [5, §5.9].)

The analysis of gaps in $\mathcal{P}(\mathbb{N})/\text{Fin}$ has played an important role in Shelah’s consistency proof of the statement that all automorphisms of the Boolean algebra $\mathcal{P}(\mathbb{N})/\text{Fin}$ are trivial ([12, §IV]). Later on, when obtaining this conclusion from PFA ([13]) one of the key parts of the proof consisted in making a given gap in $\mathcal{P}(\mathbb{N})/\text{Fin}$ indestructible. Although the gaps occurring in this proof were nonlinear, the connection with Kunen’s result is evident (it is worth noting that ‘freezing’ the gap in both cases depends on the same technique; see also [1]).

After the author has extended Shelah’s result to all quotients over analytic \mathcal{P} -ideals ([5, §3.3]) it was natural to expect that Kunen’s and Todorćević’s results will soon be extended to all quotients over analytic \mathcal{P} -ideals. It therefore came as a surprise when in January 1998 (see [5, Theorem 5.10.2]) the author has constructed an analytic Hausdorff gap \mathcal{A}, \mathcal{B} in a quotient over a certain F \mathcal{P} -ideal \mathcal{I} . (Shortly after, this object was used to construct a topological space with interesting cellularity properties, see [10].) Such a gap cannot be made indestructible, since by a result of Todorćević increasing the additivity of Lebesgue measure separates such gaps (see the proof of Lemma 7).

The proof of [5, Theorem 5.10.2] proceeded by constructing suitable families \mathcal{A}, \mathcal{B} and then dening $\mathcal{I} = \mathcal{A} \cap \mathcal{B}$. Therefore we had no control over the choice of \mathcal{I} ,

and it was rather unclear whether such a gap can be constructed in a quotient over some more natural analytic P-ideal. In [5, Question 5.13.9] and [5, Question 5.13.7] it was asked whether the quotient over the summable ideal $\mathcal{I}_{1/n} = \{A : \sum_{n \in A} 1/n < \infty\}$ satisfies some natural separation principle or has analytic Hausdorff gaps. The following implies that there are analytic Hausdorff gaps in the quotient over $\mathcal{I}_{1/n}$.

THEOREM 1. *Let \mathcal{I} be any F -P-ideal. Then there is an analytic Hausdorff gap in its quotient if and only if \mathcal{I} is not generated by a single set over Fin .*

However, it turns out that quotients over all analytic P-ideals satisfy a separation principle that implies Todorćević's separation principle in the case of Fin . This result will appear in [4].

Theorem 1 completely describes the class of all F -P-ideals that allow an analytic Hausdorff gap in their quotient, and hence gives a partial solution to [5, Problem 5.13.5], that asked for a complete description of all analytic ideals that allow an analytic Hausdorff gap in their quotient.

The rest of this note is devoted to the proof of Theorem 1. Although this paper is self-contained, familiarity with parts of [5, Chapter 1 and Chapter 5] would be useful, and we will refer to [5] for proofs of some of the results used here.

If \mathcal{A} and \mathcal{B} form an analytic Hausdorff gap in $\mathcal{P}(\mathbb{N})/\mathcal{I}$, and \mathcal{I} is an analytic P-ideal, then \mathcal{A} and \mathcal{B} are analytic P-ideals as well, and $\mathcal{A} \cap \mathcal{B} = \mathcal{I}$. By a result of Solecki (see [14]) every analytic P-ideal \mathcal{I} is of the form

$$\mathcal{I} = \text{Exh}(\mu) = \left\{ A \subseteq \mathbb{N} : \lim_{n \rightarrow \infty} \mu(A \setminus [1, n]) = 0 \right\}$$

for some lower semicontinuous submeasure μ on \mathbb{N} . In particular, every such ideal is F . Moreover, \mathcal{I} is F if and only if μ can be chosen to be *exhaustive*, i.e., so that the ideal $\text{Fin}(\mu) = \{A : \mu(A) < \infty\}$ coincides with $\text{Exh}(\mu)$. Hence the existence of an analytic Hausdorff gap in a quotient over an analytic P-ideal is equivalent to the existence of a triple of lower semicontinuous submeasures with certain properties (see [5, Proposition 5.10.1]).

The following lemma gives relatively general conditions on a lower semicontinuous submeasure μ implying the existence of an analytic Hausdorff gap in the quotient over $\text{Exh}(\mu)$.

LEMMA 2. *Assume μ is a lower semicontinuous submeasure and there are J_n , $k_n \in \mathbb{N}$, and I_{ni} ($i \leq k_n$) for $n \in \mathbb{N}$ such that*

- (1) $J_n \subseteq \mathbb{N}$ are pairwise disjoint,
- (2) $\max_{k \in J_n} \mu(\{k\}) < 2^{-n}$,
- (3) I_{ni} ($i \leq k_n$) are pairwise disjoint subsets of J_n ,
- (4) $\mu(I_{ni}) = 1$ for all $i \leq k_n$, and
- (5) $(\forall A \subseteq J_n)((\forall i \leq k_n) (\mu(A \cap I_{ni}) > 1/(n+1)^2) \rightarrow (\mu(A \cap J_n) = 1))$.

Then there is an analytic Hausdorff gap in the quotient over $\text{Exh}(\mu)$.

PROOF. For $A \subseteq \mathbb{N}$ and $n \in \mathbb{N}$ define submeasures

$$\begin{aligned} \mu_n(A) &= |\{j \in \mathbb{N} : A \cap I_{nj} \neq \emptyset\}| \\ \mu(A) &= \sup_{j \leq k_n} \mu_n(A \cap I_{nj}). \end{aligned}$$

Since both values $\nu_n(A)$ and $\nu_n(B)$ depend only on a finite set $A \cap J_n$, both submeasures

$$\begin{aligned} \nu(A) &= \sum_{n=1}^{\infty} \frac{\nu_n(A)}{n}, \\ \nu(B) &= \sup_{j \in \mathbb{N}} \nu_j(B) \end{aligned}$$

are lower semicontinuous. Let $J = \bigcup_{n=1}^{\infty} J_n$; then

$$\begin{aligned} \mathcal{A} &= \text{Exh}(\nu) \cap \mathcal{P}(J), \\ \mathcal{B} &= \text{Exh}(\nu) \cap \mathcal{P}(J) \end{aligned}$$

are analytic P-ideals. In particular, $\mathcal{A}/\text{Exh}(\nu)$ and $\mathcal{B}/\text{Exh}(\nu)$ are σ -directed. To prove that \mathcal{A}, \mathcal{B} form a Hausdorff gap in $\mathcal{P}(\mathbb{N})/\text{Exh}(\nu)$, we need only prove that \mathcal{A} and \mathcal{B} are $\text{Exh}(\nu)$ -orthogonal and that they are not separated by a single set over $\text{Exh}(\nu)$.

CLAIM 1. \mathcal{A} and \mathcal{B} are $\text{Exh}(\nu)$ -orthogonal.

PROOF. Pick $A, B \in J$. Then $\nu(A \cap B \cap J_n) \leq \nu_n(A) + \nu_n(B)$, thus

$$\nu(A \cap B) \leq \sum_{n=1}^{\infty} \nu_n(A \cap B \cap J_n) \leq \sum_{n=1}^{\infty} \frac{\nu_n(A) + \nu_n(B)}{n} = \nu(A) + \nu(B).$$

If $A \in \mathcal{A}$ and $B \in \mathcal{B}$, then $\nu(A) < \infty$ and $\lim_{m \rightarrow \infty} \nu(B \setminus \bigcup_{n=1}^m J_n) = 0$, thus by the above $\lim_{l \rightarrow \infty} \nu((A \cap B) \setminus [1, l]) = 0$, and $A \cap B \in \text{Exh}(\nu)$, as required. \square

CLAIM 2. \mathcal{A} and \mathcal{B} are not separated over $\text{Exh}(\nu)$.

PROOF. Assume the contrary, that for some $C \subseteq \mathbb{N}$ we have $A \setminus C \in \text{Exh}(\nu)$ and $B \cap C \in \text{Exh}(\nu)$ for all $A \in \mathcal{A}$ and all $B \in \mathcal{B}$. We claim that

$$(6) \quad \lim_{n \rightarrow \infty} \sup_{m \leq n, j \leq k_m} \nu(I_{mj} \setminus C) = 0.$$

Otherwise, we may find an infinite $X \subseteq \mathbb{N}$, $\varepsilon > 0$, and a ‘choice function’ $f \in \prod_{n \in X} [1, k_n]$ such that

$$\nu(I_{nf(n)} \setminus C) > \varepsilon$$

for all $n \in X$. We may furthermore shrink X so that $\sum_{n \in X} 1/n < \infty$. Let $A = \bigcup_{n \in X} I_{nf(n)}$; then $\nu(A) = \sum_{n \in X} \varepsilon/n < \infty$, thus $A \in \mathcal{A}$. Note that $A \cap C = \emptyset$. However, for $n \in X$ we have $\nu(A \cap J_n) = \nu(A \cap I_{nf(n)}) > \varepsilon$, therefore $A \notin \text{Exh}(\nu)$, contradicting the assumption on C .

By (6) for all but finitely many n we have $\sup_{j \leq k_n} \nu(I_{nj} \setminus C) < 1/2$, and therefore

$$(7) \quad \nu(I_{nj} \cap C) > 1/2 \text{ for all but finitely many } n \text{ and all } j \leq k_n.$$

Let Y be the set of such n . We shall now find $B_n \subseteq J_n \cap C$ ($n \in Y$) so that for all $n \in Y$ we have

$$\begin{aligned} (8) \quad \nu(B_n) &< 1, \\ (9) \quad \nu_n(B_n) &< 1/n^2. \end{aligned}$$

Once these sets are constructed, let $B = \bigcup_{n \in Y} B_n$. Then $B \subseteq C$ and $\nu_n(B) < 1/n$, thus $B \in \mathcal{B}$, yet $B \notin \text{Exh}(\nu)$ by (8). Since this contradicts the assumption that C separates \mathcal{A} and \mathcal{B} , it will suffice to describe the construction of B_n ($n \in Y$).

Fix $n \in Y$. By (7) and (2) we may find $B_n \subseteq J_n \cap C$ such that

$$(10) \quad 1/n^2 < \nu_n(B_n \cap I_{nj}) < 1/n^2.$$

for all $j \leq k_n$, thus (9) holds. By (5), $(B_n \cap J_n) \leq 1$, thus (8) holds as well. This completes the proof. \square

Claim 1 and Claim 2 complete the proof of lemma. \square

LEMMA 3. *Assume that \mathcal{I} is an exhaustive lower semicontinuous submeasure such that $\lim_{n \rightarrow \infty} (\{n\}) = 0$ and $\text{Exh}(\mathcal{I})$ is a proper ideal. Then \mathcal{I} satisfies the conditions of Lemma 2.*

PROOF. Since $\text{Fin}(\mathcal{I}) = \text{Exh}(\mathcal{I})$ is a proper ideal, we have $(\mathbb{N}) = \infty$. We describe the recursive construction of J_n ($n \in \mathbb{N}$). Suppose J_i ($i \leq n$) have already been constructed. Since $\lim_{k \rightarrow \infty} (\{k\}) = 0$ and \mathcal{I} is lower semicontinuous, we may find m such that

$$\bigcup_{i=1}^{n-1} J_i \leq [1, m] \quad \text{and} \quad \sup_{j \leq m} (\{j\}) < 2^{-n}.$$

We will have $\min(J_n) \leq m$, assuring (1) and (2). Since $(\mathbb{N}) = \infty$ and \mathcal{I} is lower semicontinuous, we can find $m = m_0 < m_1 < m_2 < \dots$ such that $1 \leq [m_i, m_{i+1})$ for all i . Let T be the family of all $B \leq [m, \infty)$ such that for some $j \in \mathbb{N}$ and all $i \leq j$ we have

$$(11) \quad 1/n^2 < \mathcal{I}(B \cap [m_i, m_{i+1})), \text{ and}$$

$$(12) \quad \mathcal{I}(B) \leq 1.$$

We claim that $T \in \mathcal{P}([m_0, m_k])$ for some $k \in \mathbb{N}$. Otherwise T is infinite, and by König's lemma there is an infinite sequence of sets $B_i \leq [m_i, m_{i+1})$ whose union satisfies (11) for all i yet has submeasure at most 1. But this contradicts the exhaustivity of \mathcal{I} . Thus $T \in \mathcal{P}([m_0, m_k])$ for some k . Let $k_n = m_{k+1}$, and $I_{ni} = [m_{i-1}, m_i)$ for $i \leq k_n$. Then clearly (3) and (4) hold. Let us check (5). If $A \leq J_n$ is such that $\mathcal{I}(A \cap I_{ni}) > 1/n^2$ for all $i \leq k_n$, then since $A \leq [m_0, m_k)$ we have $A \notin T$, and therefore $\mathcal{I}(A) > 1$, as required.

This describes the recursive construction of sets satisfying the assumptions of Lemma 2. \square

I do not know whether the assumption that \mathcal{I} is exhaustive is necessary in Lemma 3, but a \mathcal{I} satisfying all assumptions of Lemma 3 with exhaustivity replaced by $(\mathbb{N}) = \infty$ and failing the conclusion of Lemma 3 would have to be rather peculiar.

An ideal \mathcal{I} is *dense* if every infinite set has an infinite subset in \mathcal{I} . It is *dense below* A (for some $A \leq \mathbb{N}$) if every infinite subset of A has an infinite subset in \mathcal{I} . Note that $\text{Exh}(\mathcal{I})$ is dense if and only if $\lim_{n \rightarrow \infty} (\{n\}) = 0$. An ideal has the *Fechet property* if it is not dense below any positive set. The following special case of a result of Todorcevic shows that it suffices to prove Theorem 1 for dense ideals.

LEMMA 4. *An F - \mathcal{P} -ideal has the Fechet property if and only if it is generated by a single set over Fin .*

PROOF. This follows immediately from [18, Theorem 2]; for a proof see [5, Corollary 1.2.11]. \square

PROOF OF THEOREM 1. The direct implication follows from [17, Corollary 1], since an ideal generated by a single set over Fin is isomorphic to Fin .

Let \mathcal{I} be an F - \mathcal{P} -ideal not generated by a single set over Fin . By Solecki's theorem, let \mathcal{I} be a lower semicontinuous exhaustive submeasure such that $\mathcal{I} =$

Exh(). By Lemma 4, there is an \mathcal{I} -positive set A such that \mathcal{I} is dense below A ; we may assume that $A = \mathbb{N}$, i.e., that \mathcal{I} is dense. This is easily equivalent to $\lim_{k \rightarrow \infty} (\{k\}) = 0$. By the exhaustivity of \mathcal{I} , $\text{add}(\mathcal{I}) = \infty$, thus by Lemma 3 the assumptions of Lemma 2 are satisfied and there is an analytic Hausdorff gap in $\mathcal{P}(\mathbb{N})/\mathcal{I}$ as required. \square

Recall that the *summable ideal* is denoted by

$$\mathcal{I}_{1/n} = \left\{ A \subseteq \mathbb{N} : \sum_{n \in A} \frac{1}{n} < \infty \right\}.$$

For some time Fin and $\mathcal{I}_{1/n}$ (and its obvious variations) were essentially the only known examples of F -P-ideals, until a number of radically different examples was found (see [5, §1.11]), but $\mathcal{I}_{1/n}$ still stands out as the simplest nontrivial F -P-ideal.

COROLLARY 5. *There is an analytic Hausdorff gap in $\mathcal{P}(\mathbb{N})/\mathcal{I}_{1/n}$.* \square

Corollary 6 below gives partial answers to [19, Problem 2], [5, Problem 5.13.16], and [5, Question 5.13.17]. Together with the Kunen–Todorćević result, it also shows that the gap-spectra of quotients over F -P-ideals other than Fin radically differ from the gap-spectrum of $\mathcal{P}(\mathbb{N})/\text{Fin}$. Recall that the ideals \mathcal{I} and \mathcal{J} are *isomorphic* if there are $A \in \mathcal{I}$, $B \in \mathcal{J}$ and a bijection h between $\mathbb{N} \setminus A$ and $\mathbb{N} \setminus B$ such that $C \in \mathcal{I}$ if and only if $h^{-1}(C) \in \mathcal{J}$. Clearly, this implies that the quotients over \mathcal{I} and \mathcal{J} are isomorphic.

COROLLARY 6. *Assume Martin’s Axiom. Then there is a $(\mathfrak{c}, \mathfrak{c})$ -gap in the quotient over every F -P-ideal not isomorphic to Fin .*

PROOF. Since Martin’s Axiom implies that $\text{add}(\mathcal{N}) = \text{cof}(\mathcal{N}) = \mathfrak{c}$, this follows immediately from Theorem 1 and Lemma 7 below. \square

Recall that $\text{add}(\mathcal{N})$ is the additivity of the Lebesgue measure and that $\text{cof}(\mathcal{N})$ is the conality of the ideal of Lebesgue null sets.

LEMMA 7. *Assume $\text{add}(\mathcal{N}) = \text{cof}(\mathcal{N})$. Then there is an $(\text{add}(\mathcal{N}), \text{add}(\mathcal{N}))$ -gap in the quotient over every analytic P-ideal whose quotient contains an analytic Hausdorff gap.*

PROOF. This proof is identical to the proof of [5, Lemma 5.6.3], but we recall it for the reader’s convenience. Let \mathcal{I} be an analytic P-ideal and let $\mathcal{A}_0, \mathcal{B}_0$ be an analytic Hausdorff gap in $\mathcal{P}(\mathbb{N})/\mathcal{I}$. Let \mathcal{A} be the ideal generated by \mathcal{A}_0 and \mathcal{I} , and let \mathcal{B} be the ideal generated by \mathcal{B}_0 and \mathcal{I} . Both \mathcal{A} and \mathcal{B} are analytic P-ideals.

By a result of Todorćević ([18, Theorem 5]), for every analytic P-ideal \mathcal{A} the partially ordered set (\mathcal{A}, \subseteq) (i.e., \mathcal{A} ordered by the inclusion modulo finite) is Tukey-reducible to Lebesgue null sets ordered by the inclusion (see also [5, Theorem 5.6.2]). In particular, every subset X of \mathcal{A} of size less than $\text{add}(\mathcal{N})$ is bounded in (\mathcal{A}, \subseteq) , and there is a $Y \in \mathcal{I}$ of size $\text{cof}(\mathcal{N})$ such that Y is \mathcal{I} -conal. Since $\text{add}(\mathcal{N}) = \text{cof}(\mathcal{N})$, we can recursively construct two chains, $\mathcal{A}' \subseteq \mathcal{A}$ and $\mathcal{B}' \subseteq \mathcal{B}$, increasing modulo Fin and each of length $\text{add}(\mathcal{N})$ such that \mathcal{A}' is conal in \mathcal{A} and \mathcal{B}' is conal in \mathcal{B} . Hence $\mathcal{A}', \mathcal{B}'$ form an $(\text{add}(\mathcal{N}), \text{add}(\mathcal{N}))$ gap in $\mathcal{P}(\mathbb{N})/\mathcal{I}$. \square

Note that Lemma 7 shows that an analytic Hausdorff gap in a quotient over an analytic P-ideal can always be filled by a ccc forcing, contrasting the Kunen’s result on the indestructibility of gaps in $\mathcal{P}(\mathbb{N})/\text{Fin}$. It should be noted that the

statement ‘ $\mathcal{P}(\mathbb{N})/\text{Fin}$ contains a $(\mathfrak{c}, \mathfrak{c})$ Hausdorff gap’ is independent from Martin’s Axiom (Kunen, [7]).

The only two analytic P-ideals (up to the isomorphism) that are known not to have analytic Hausdorff gaps in their quotients are Fin (by [19, Corollary 1]) and $\emptyset \text{---} \text{Fin}$ (by [5, Proposition 5.3.2] and Lemma 7). By a result of Todorćević (a special case of which is Lemma 4), these are also the only two analytic P-ideals that have the \mathfrak{F} chet property (see [5, Corollary 1.2.11]).

QUESTION 8. (a) *Assume \mathcal{I} is an analytic P-ideal that does not have the \mathfrak{F} chet property. Is there an analytic Hausdorff gap in its quotient?*

(b) *Assume \mathcal{I} is an F -ideal that does not have the \mathfrak{F} chet property. Is there an analytic Hausdorff gap in its quotient?*

(Note that Fin is not the only F -ideal with the \mathfrak{F} chet property; e.g., the ideal Ib of [9, §1.4] generated by the branches of the tree $\mathbb{N}^{<\mathbb{N}}$ has the \mathfrak{F} chet property.) Let us explicitly state a special case of Question 8(a).

QUESTION 9. *Assume a dense analytic P-ideal is equal to $\text{Exh}(\mu)$ for a lower semicontinuous submeasure μ such that $\mu(\mathbb{N}) = \infty$. Is there an analytic Hausdorff gap in its quotient?*

Lemma 2 can be used to give a positive answer to this question for many ‘natural’ submeasures μ . While a positive solution to Question 9 may not require much beyond what is given in the proof of Theorem 1, the following question most likely requires a different approach.

QUESTION 10. *Is there an analytic Hausdorff gap in the quotient over the ideal of asymptotic density zero sets, $\mathcal{Z}_0 = \{A : \limsup_n |A \cap n|/n = 0\}$?*

The existence of analytic Hausdorff gaps does not immediately give new linear gaps, unless MA or a similar axiom is assumed (see Lemma 7; to see why linear Hausdorff gaps are of some interest, see e.g., [10]). The following is a stronger version of [5, Conjecture 5.13.4].

CONJECTURE 11. *There is an $\langle \text{add}(\mathcal{N}), \text{add}(\mathcal{N}) \rangle$ -gap in the quotient over every F -P-ideal that is not generated by a single set over Fin .*

A confirmation of Conjecture 11 would give a positive answer to [19, Problem 3], where it was asked whether some standard cardinal invariants of the continuum other than ω, ω_1 and the bounding number for $\mathbb{N}^{\mathbb{N}}/\text{Fin}$ naturally occur in the (linear) gap spectra of some analytic quotient.

At the end, let us point out that analytic Hausdorff gaps exist in quotients over analytic ideals that are not P-ideals. For example, the quotient over the following ideal (otp stands for the order-type of a subset of an ordinal)

$$\mathcal{I}_{\omega^2} = \{A \subseteq \omega^2 : \text{otp}(A) < \omega^2\}$$

(also known as $\text{Fin} \text{---} \text{Fin}$) has such a gap (see [5, Proposition 5.8.3]).

References

- [1] U. Abraham, M. Rubin, and S. Shelah. On the consistency of some partition theorems for continuous colorings, and the structure of \aleph_1 -dense real order types. *Annals of Pure and Applied Logic*, 29:123–206, 1985.
- [2] M. Bekkali. *Topics in Set Theory*, volume 1476 of *Lecture Notes in Mathematics*. Springer-Verlag, 1991.

- [3] H.G. Dales and W.H. Woodin. *An Introduction to Independence for Analysts*, volume 115 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, 1987.
- [4] I. Farah. Luzin gaps, separation principles and preservation of gaps. in preparation.
- [5] I. Farah. *Analytic quotients: theory of liftings for quotients over analytic ideals on the integers*. Number 702 in *Memoirs of the American Mathematical Society*, vol. 148. 2000. 177 pp.
- [6] F. Hausdor. Summen von \aleph_1 Mengen. *Fundamenta Mathematicae*, 26:241–255, 1936.
- [7] K. Kunen. $\langle \cdot, \cdot \rangle$ -gaps under MA. preprint, 1976.
- [8] A.R.D. Mathias. A remark on rare lters. In A. Hajnal et al., editors, *Infinite and finite sets, Vol. III*, volume 10 of *Coll. Math. Soc. János Bolyai*, pages 1095–1097. North Holland, 1975.
- [9] K. Mazur. F -ideals and $\omega_1\omega_1$ -gaps in the Boolean algebra $\mathcal{P}(\omega)/I$. *Fundamenta Mathematicae*, 138:103–111, 1991.
- [10] J.T. Moore. Linearly bared Souslinean space under MA. *Topology Proceedings*, 24, Spring 1999.
- [11] M. Scheepers. Gaps in ${}^\omega\omega$. In *Set theory of the reals*, volume 6 of *Israel Mathematical Conference Proceedings*, pages 439–561. 1993.
- [12] S. Shelah. *Proper Forcing*. Lecture Notes in Mathematics 940. Springer, 1982.
- [13] S. Shelah and J. Steprans. PFA implies all automorphisms are trivial. *Proceedings of the American Mathematical Society*, 104:1220–1225, 1988.
- [14] S. Solecki. Analytic ideals and their applications. *Annals of Pure and Applied Logic*, 99:51–72, 1999.
- [15] R. Solovay. A model of set theory in which every set of reals is Lebesgue measurable. *Annals of Mathematics*, 92:1–56, 1970.
- [16] S. Todorcevic. *Partition Problems in Topology*, volume 84 of *Contemporary mathematics*. American Mathematical Society, Providence, Rhode Island, 1989.
- [17] S. Todorcevic. Analytic gaps. *Fundamenta Mathematicae*, 150:55–67, 1996.
- [18] S. Todorcevic. Denable ideals and gaps in their quotients. In C.A. DiPrisco et al., editors, *Set Theory: Techniques and Applications*, pages 213–226. Kluwer Academic Press, 1997.
- [19] S. Todorcevic. Gaps in analytic quotients. *Fundamenta Mathematicae*, 156:85–97, 1998.

DEPARTMENT OF MATHEMATICS, CUNY, COLLEGE OF STATEN ISLAND AND GRADUATE CENTER,
 AND MATEMATICKI INSTITUT, KNEZA MIHAILA 35, BELGRADE
E-mail address: ifarah@gc.cuny.edu
URL: <http://www.math.csi.cuny.edu/farah>