

ISSN:1120-6330

ATTI DELLA ACCADEMIA NAZIONALE DEI LINCEI

ANNO CCCXCII - 1995

CLASSE DI SCIENZE FISICHE, MATEMATICHE E NATURALI

RENDICONTI LINCEI

MATEMATICA E APPLICAZIONI

SERIE IX - VOLUME VI - FASCICOLO 4

ESTRATTO



Comitato consultivo

A. Ambrosetti (Pisa), E. Arbarello (Roma), V. Arnold (Mosca), G. Capriz (Pisa),
C. Cercignani (Milano), M. Cornalba (Pavia), G. Fichera (*Direttore*, Roma),
D. Galletto (Torino), E. Gatti (Milano), G. Grioli (Padova), W. Hayman (Londra),
G. Letta (Pisa), E. Magenes (Pavia), G. Maier (Milano), E. Marchi (Genova),
L. Nirenberg (New York), O. Oleinik (Mosca), C. Procesi (Roma), E. Vesentini
(Pisa), G. Zappa (Firenze)

Teoria dei gruppi. — *On absolutely-nilpotent of class k groups.* Nota (*) di PATRIZIA LONGOBARDI, TRUEMAN MACHENRY, MERCEDE MAJ e JAMES WIEGOLD, presentata dal Socio G. Zappa.

ABSTRACT. — A group G in a variety \mathfrak{V} is said to be absolutely- \mathfrak{V} , and we write $G \in A\mathfrak{V}$, if central extensions by G are again in \mathfrak{V} . Absolutely-abelian groups have been classified by F. R. Beyl. In this paper we concentrate upon the class $A\mathcal{N}_k$ of absolutely-nilpotent of class k groups. We prove some closure properties of the class $A\mathcal{N}_k$ and we show that every nilpotent of class k group can be embedded in an $A\mathcal{N}_k$ -group. We describe all metacyclic $A\mathcal{N}_k$ -groups and we characterize 2-generator and infinite 3-generator $A\mathcal{N}_2$ -groups. Finally we study extensions $1 \rightarrow N \rightarrow H \rightarrow G \rightarrow 1$, with $N \leq \zeta_n(H)$, the n -centre of H , with $n > 1$.

KEY WORDS: Variety; Central extension; Nilpotent group.

RIASSUNTO. — *Gruppi assolutamente-nilpotenti di classe k .* Un gruppo G in una varietà \mathfrak{V} vien detto assolutamente- \mathfrak{V} (e si scrive $G \in A\mathfrak{V}$) se ogni estensione centrale mediante G appartiene ancora a \mathfrak{V} . I gruppi assolutamente-abeliani sono stati caratterizzati da F. R. Beyl. In questa *Nota* si studiano i gruppi assolutamente-nilpotenti di classe k . Si provano alcune proprietà di chiusura della classe $A\mathcal{N}_k$, e si mostra che ogni gruppo nilpotente di classe k si può immergere in un $A\mathcal{N}_k$ -gruppo. Si descrivono i gruppi metaciclici assolutamente-nilpotenti di classe k ed i gruppi 2-generati e quelli infiniti 3-generati nella classe $A\mathcal{N}_2$. Infine si esaminano estensioni $1 \rightarrow N \rightarrow H \rightarrow G \rightarrow 1$, con $N \leq \zeta_n(H)$, l' n -esimo centro di H .

1. INTRODUCTION

A group G in a variety \mathfrak{V} is said to be *absolutely- \mathfrak{V}* if central extensions by G are again in \mathfrak{V} . We denote the class of absolutely- \mathfrak{V} groups by $A\mathfrak{V}$.

Special cases of such groups have been considered by several authors, *e.g.* Varadarajan [9], Evens [5] and Beyl [1-3].

In this paper we concentrate upon the class $A\mathcal{N}_k$ of absolutely-nilpotent of class k groups. In [1] Beyl has classified all absolutely abelian groups; they are just those abelian groups having trivial multipliers. Conditions sufficient to ensure that groups in nilpotent varieties are absolute are studied in Passi and Vermani [7].

In sect. 2 of this paper we collect some general results about $A\mathcal{N}_k$ -groups. Obviously the class $A\mathcal{N}_k$ is not a variety, for example, it is not necessarily subgroup-closed, but it does have some interesting closure properties: for instance, it is closed under some nilpotent products (see 2.1). We do not know if $A\mathcal{N}_k$ is closed under Cartesian products, the best we can say here is that Cartesian powers of finite $A\mathcal{N}_k$ -groups are $A\mathcal{N}_k$ -groups (see 2.3). We have not been able to recognize if epimorphic images of an $A\mathcal{N}_k$ -group G are always $A\mathcal{N}_k$, we have only proved that this is true if G is finitely generated (see 2.4).

Every nilpotent n -generator group of class k can be embedded in a $2n$ -generator $A\mathcal{N}_k$ -group (see 2.2), in particular the class of n -generator $A\mathcal{N}_k$ -groups is not a small class. In sect. 3, 4 and 5 we concentrate upon \mathcal{N}_k -groups with 2 or 3 generators.

(*) Pervenuta all'Accademia il 5 luglio 1995.

We describe all metacyclic $A\mathcal{N}_k$ -groups (see [4], for $k = 2$). Moreover we characterize the 2-generator and the infinite 3-generator class 2 groups which are absolute-there are no infinite 2-generator non abelian $A\mathcal{N}_2$ -groups, and, in a sense, very few infinite 3-generator $A\mathcal{N}_2$ -groups.

Finally, in sect. 6 we study extensions $1 \rightarrow N \rightarrow H \rightarrow G \rightarrow 1$ with $N \leq \xi_n(H)$, $n > 1$.

Notation is as in [8].

If $n \geq 1$ is an integer, x, y elements of a group G , H a subgroup of G , we put $[x, {}_n y] = [x, y, \dots, y]$, $[H, {}_n G] = [H, G, \dots, G]$.

If $F = \langle x, y \rangle$ is a free group, we write

$$d_{i,n} = [x, y, x, \dots, x, y, \dots, y]_{i, n-i}, \quad \text{for any } n \geq 0, 0 \leq i \leq n.$$

Obviously $[d_{i,n}, x] \equiv d_{i+1, n+1} \pmod{F''}$ and $[d_{i,n}, y] \equiv d_{i, n+1} \pmod{F''}$.

Moreover the set $\{d_{i,n} \mid n \in \mathbb{N}_0, i \in \{0, 1, \dots, n\}\}$ is a basis for $F' \pmod{F''}$.

The following result will be used frequently:

1.1. (see [7, Theorem 5.1]) Let \mathfrak{V} be a variety of exponent 0. Then $G \in A\mathfrak{V}$ if and only if $\mathfrak{V}(F) \leq [R, F]$ for all free presentations $1 \rightarrow R \rightarrow F \rightarrow G \rightarrow 1$.

2. GENERAL RESULTS

It is easy to see that $A\mathcal{N}_k$ is closed under restricted direct products. In fact the result holds even for generalized nilpotent products.

2.1. Let $\{A_i : i \in I\}$ be any set of class k groups.

If $A_i \in A\mathcal{N}_k$ for any $i \in I$ and $l < k$, then the l -th nilpotent product of the A_i is in $A\mathcal{N}_k$.

PROOF. Let G be such a group, and let $1 \rightarrow R \rightarrow F \xrightarrow{\pi} G \rightarrow 1$ be a free presentation of G . By 1.1 we have to prove that $\gamma_{k+1} F \leq [R, F]$.

Write $X_i = \pi^{-1}(A_i)$, for any $i \in I$; obviously it suffices to prove that $[y_1, y_2, \dots, y_{k+1}] \in [R, F]$ for any $y_1, y_2, \dots, y_{k+1} \in \bigcup_{i \in I} X_i$. If there exist $r, s \leq k, r \neq s$, such that $y_b \in X_r, y_j \in X_s$, for some $b, j \leq k$, then $[y_1, y_2, \dots, y_k] \in [X_r, X_s]^F \cap \gamma_k F \leq R$ and $[y_1, y_2, \dots, y_k, y_{k+1}] \in [R, F]$. If y_1, y_2, \dots, y_k are in one and the same X_i , and $y_{k+1} \in X_j$, then $[y_1, y_2, \dots, y_k, y_{k+1}] \in [R \cap \langle X_i \rangle, \langle X_j \rangle] \subseteq [R, F]$, since $A_i \in A\mathcal{N}_k$. Now assume that $y_1, \dots, y_k \in X_r$ and $y_{k+1} \in X_s$, for suitable $r \neq s$. Then $[y_1, y_2, \dots, y_k, y_{k+1}] \in [\gamma_k X_r, X_s]$. But, for any $j < k$, $[X_s, X_r, \dots, X_r, \gamma_{k-j-1} X_r] \leq \leq \gamma_k F \cap [X_s, X_r]^F \leq R$ and $[X_s, X_r, \dots, X_r, \gamma_{k-j-1} X_r, X_r] \leq [R, F]$.

Hence, by induction on i , using the three subgroups lemma, it is easy to verify that $[\gamma_i X_r, [X_s, X_r, \dots, X_r]] \leq [R, F]$ for $1 \leq i \leq k$. Hence with $i = k$ we have $[\gamma_k X_r, X_s] \leq [R, F]$, as required. \blacksquare

For some nilpotent products we can drop the condition that the factors be absolute.

2.2. Let $\{A_i: i \in I\}$ be any set of class k groups such that $\gamma_k A_i \cong \gamma_k A_j$, $i, j \in I$. Then, with $l < k$, the l -th nilpotent product of all A_i with $\gamma_k A_i$ amalgamating is in $A\mathcal{N}_k$.

PROOF. Similar to the proof of 2.1. ■

For Cartesian products we have:

2.3. Let A be a finite $A\mathcal{N}_k$ -group. Then every Cartesian power of A is in $A\mathcal{N}_k$.

PROOF. Let A^I be the Cartesian power of A with index set I , and G a group having a central subgroup N such that $G/N \cong A^I$.

To show that G is in \mathcal{N}_k , it is enough to show that every finitely generated subgroup of G is in \mathcal{N}_k .

Thus, let H be a finitely generated subgroup of G . Then HN/N is a finitely generated subgroup of A^I , and the finiteness of A now implies that HN/N is a subgroup of the direct product X of finitely many groups isomorphic to A , namely of diagonals of powers A^J with $J \subseteq I$. (The reader will recognize the genesis of this type of argument in B. H. Neumann's proof [6] that Cartesian products of finite groups are locally finite). Note next that X is in $A\mathcal{N}_k$. Thus, if K is the subgroup of G such that $K/N = X$, it follows that K is in \mathcal{N}_k , and thus H is in \mathcal{N}_k since $K \geq H$.

Therefore G is in \mathcal{N}_k and A^I is in $A\mathcal{N}_k$, as required. ■

It would be nice to know if epimorphic images of $A\mathcal{N}_k$ -groups are always $A\mathcal{N}_k$. We have not been able to confirm or deny this. For finitely generated groups, we have:

2.4. If G is a finitely generated $A\mathcal{N}_k$ -group and N is a normal subgroup of G , then G/N is in $A\mathcal{N}_k$.

PROOF. Let G be a finitely generated nilpotent group. Then $M(G)$ is finitely generated and so *Hopf*. Write $\mathfrak{V} = \mathcal{N}_k$.

Obviously it suffices to show that G is in $A\mathfrak{V}$ if and only if $\mathfrak{V}(F) \leq [R, F]$ for some free presentation $1 \rightarrow R \rightarrow F \rightarrow G \rightarrow 1$ of G . For, let $1 \rightarrow R_1 \rightarrow F_1 \rightarrow G \rightarrow 1$ any free presentation of G . Then $(R_1 \cap F_1') / ([R_1, F_1] V(F_1)) \cong M_V(G) \cong M(G) \cong (R_1 \cap F_1') / [R_1, F_1]$, so that $\mathfrak{V}(F_1) \leq [R_1, F_1]$. Then the result follows from 1.1. ■

We end this section with the following theorem, which in some sense reduces the study of finitely generated $A\mathcal{N}_k$ -groups to the cases of torsion-free groups and finite groups (see also [1, Theorem 2.1]).

2.5. Let $G \in \mathcal{A}\mathcal{N}_k$. If G is finitely generated, then G can be embedded as a subgroup of finite index in an $\mathcal{A}\mathcal{N}_k$ -group which is the direct product of a finite $\mathcal{A}\mathcal{N}_k$ -group and a finitely generated torsion-free $\mathcal{A}\mathcal{N}_k$ -group.

PROOF. Let τG be the torsion subgroup of G . Then τG is finite and $G/\tau G$ is a finitely generated torsion-free $\mathcal{A}\mathcal{N}_k$ -group by 2.4. Since G is residually finite, there exists a normal subgroup N in G such that $N \cap \tau G = 1$ and G/N is a finite group in $\mathcal{A}\mathcal{N}_k$, again by 2.4. Then $\chi: x \in G \mapsto (xN, x\tau G) \in G/N \times G/\tau G$ is an embedding. By 2.1 $H = G/N \times G/\tau G \in \mathcal{A}\mathcal{N}_k$ if $k > 1$. Moreover G^χ has finite index in H since N^χ has finite index in $G/\tau G$. If $k = 1$, the result is trivial. ■

3. METACYCLIC $\mathcal{A}\mathcal{N}_k$ -GROUPS

THEOREM 3.1. Let G be a metacyclic p -group of class k . Then

(A) $G \in \mathcal{A}\mathcal{N}_k$ if and only if one of the following holds:

(i) $G = \langle x, y \mid |x| = p^a, x^{p^b} = y^{p^c}, [x, y] = x^{p^d} \rangle$, with $(p, s) = 1$, $a \leq b + d$, $b \leq c$, and $b \leq (k-1)d$, $d > 0$;

(ii) $p = 2$, $k > 2$ and $G = \langle x, y \mid |x| = 2^k, y^{2^{k-m-1}} = x^{2^k} = 1, x^y = x^{-1+2^m} \rangle$, with $0 < m < k-1$;

(iii) $p = 2$, $k > 2$ and $G = \langle x, y \mid |x| = 2^k, x^{2^{k-1}} = y^{2^c}, x^y = x^{-1+2^m} \rangle$, with $s \neq 0(2)$, $0 < m < k-1$, $k-1-m \leq c \leq k-1$;

(iv) $p = 2$, $k > 2$ and $G = \langle x, y \mid |x| = 2^k, x^{2^{k-1}} = y^{2^b}, x^y = x^{-1} \rangle$, with $0 < b < k-1$.

(B) For any $n \geq 2$ there exists a group H , with a normal subgroup $M \leq \zeta_n H$ such that $H/M \cong G$ and $clH \geq n + k - 1$.

PROOF. Since G is metacyclic, $G = \langle x, y \rangle$, with $|x| = p^a$, $x^{p^b} = y^{p^c}$, $[x, y] = x^{p^d}$, $a \leq b + d$, $(p, s) = 1$, $d \geq 1$.

Let $1 \rightarrow R \rightarrow F \rightarrow G \rightarrow 1$ be a free presentation of G , with $F = \langle \bar{x}, \bar{y} \rangle$.

Assume first that $p \neq 2$. Then $(xy)^{p^{a-1}} = x^{p^{a-1}}y^{p^{a-1}}$, and we may suppose $|x| \leq |y|$ and $b \leq c$ (replacing eventually y by xy). Then $[R, F] = \langle d_{i,n}, d_{0,b}^{p^b}, d_{0,r}, d_{0,m}^{-1}d_{0,0}^{p^{md_s m}} \rangle F''$, with $1 \leq n$, $1 \leq i \leq n$, $1 \leq m \leq k-1$, $0 \leq b \leq k-1$, $r \geq s$. From $\gamma_{k+1}F \leq [R, F]$ it follows that $d_{0,k-1} \in [R, F]$ and

$$d_{0,k-1} = (d_{0,k-1}^{-1}d_{0,0}^{p^{(k-1)d_s k-1}})^{\beta} d_{0,0}^{p^b \alpha} (d_{0,k-1})^{p^b \gamma}$$

since the $d_{b,l}$ are independent mod F'' .

Hence $\beta - p^b \gamma - 1 = 0$, $p^{(k-1)d_s k-1} \beta + p^b \alpha = 0$ and that happens if and only if $b \leq (k-1)d$.

Therefore if G is in $\mathcal{A}\mathcal{N}_k$ then $b \leq (k-1)d$.

Conversely it is easy to see that G is in $\mathcal{A}\mathcal{N}_k$ if (i) holds.

Furthermore, for any $l \geq 2$ we have

$$[N, {}_l F] \leq F'' \langle d_{i,n}, d_{0,b+l}^{p^b}, d_{0,r+l}, d_{0,m+l}^{-1} d_{0,l}^{p^{md_s m}} \rangle$$

$1 \leq n, k \leq r, 1 \leq m \leq k-1, 1 \leq i$.

From $\gamma_{k+l-2} F = [N, {}_{l-1} F]$ it follows $d_{0,k-2+l-1} \in [N, {}_{l-1} F]$ and

$$d_{0,k+l-3} = (d_{0,k-2+(l-1)}^{-1} d_{0,l-1}^{(k-2)d_s k-2})^\beta (d_{0,l}^{p^b})^\alpha (d_{0,k+l-2}^{p^b})^\gamma$$

and from that $b \leq (k-2)d$. Thus $a \leq b+d \leq (k-1)d$ and G is of class $\leq k-1$, a contradiction.

Then, with $T = [N, {}_{l-1} F]$, the group $H = F/T$ has class $\geq s+l-1$ and $N/T \leq \zeta_l H$, hence (\mathcal{B}) holds.

Now assume that $p = 2$.

If $|x| \leq |y|$, arguing as before we can prove that G has the structure in (i) and that (\mathcal{B}) holds. Now let $|x| > |y|$; then we can assume $x^y = x^{1+2^t}$, with t odd (if $[x, y] = x^{4^s}$, then $(yx)^{2^{s-1}} = x^{2^{s-1}} \neq 1$ and, replacing y by xy , we get $|x| \leq |y|$).

Then $[x, {}_i y] = x^{2^i}$, for every i , and $|x| = 2^k$ since G has class k , also $b \geq k-1$.

First suppose that $t \neq -1$; then $t = -1 + 2^m s$ with $s \neq 0(2)$. By induction on b it is easy to prove that, for every $b \in \mathbb{N}$, we have $[x, y^{2^b}] = [x, y]^{2^{b+m}\beta}$, $\beta \neq 0(2)$. Thus $c \geq k-m-1$, and $[N, F] = \langle d_{i,n}, d_{0,b}^{2^{c+m}}, d_{0,b}^{2^b}, d_{0,r}, d_{0,l}^{-1} d_{0,0}^{2^{2^l l}} \rangle F''$ with $n \in \mathbb{N}$, $1 \leq i \leq n, 0 \leq b \leq k-1, r \geq k, 1 \leq l \leq k-1$. If $b > c+m$ and $G \in \mathcal{A}\mathcal{X}_k$ then $c+m \leq k-1$, and so $c+m = k-1, c = k-m-1, b = k$ and (ii) holds. If $b \leq c+m$ and $G \in \mathcal{A}\mathcal{X}_k$, then $b \leq k-1$, thus $b = k-1$ and (iii) holds.

As in the previous case we prove that (\mathcal{B}) holds and that $G \in \mathcal{A}\mathcal{X}_k$ if either (ii) or (iii) hold.

Finally suppose $x^y = x^{-1}$. Then, if $b = k-1$, we get

$$G = \langle x, y | x^{2^k} = 1, x^{2^{k-1}} = y^{2^b}, x^y = x^{-1} \rangle, \quad 1 \leq b < k-1,$$

and $[N, F] = \langle d_{i,n}, d_{0,b}^{2^{k-1}}, d_{0,r}, d_{0,m} d_{0,0}^{2^{2^m}} \rangle F''$, with $n \in \mathbb{N}, 1 \leq i \leq n, 0 \leq b \leq k-1, r \geq k, 1 \leq m \leq k-1$. Thus $d_{0,k-1} \in [N, F]$ and arguing as in the previous case $d_{0,k+l-3} \notin [N, {}_{l-1} F]$. If $b = k$, then $G = \langle x \rangle \rtimes \langle y \rangle$, with $|x| = 2^k, x^y = x^{-1}, |y| = 2^b, 0 < b < k$. Then, with $G_n = \langle a \rangle \rtimes \langle b \rangle, |a| = 2^{k+b}, a^b = a^{-1}, |b| = 2^b$, we have $a^{2^k} \in \zeta_n G_n$, and G_n has class $k+n$. From $G_n / \langle a^{2^k} \rangle \cong G$ it follows that G is not in $\mathcal{A}\mathcal{X}_k$ and that (\mathcal{B}) holds. ■

4. 2-GENERATOR $\mathcal{A}\mathcal{X}_2$ -GROUPS

We give a complete description of 2-generator $\mathcal{A}\mathcal{X}_2$ -groups.

We start with an easy Lemma:

LEMMA 4.1. Let G be a 2-generator finite p -group and $1 \rightarrow R \rightarrow F \rightarrow G \rightarrow 1$ a free presentation of G , with F 2-generator. Then either G is metacyclic or $R \leq F^p [F', F]$ and, if $p = 2, R \leq \langle x^2, y^2 \rangle (F')^2 [F', F]$, with $F = \langle x, y \rangle$.

PROOF. There exist free generators x, y of F such that

$$R = \langle x^\alpha [x, y]^\gamma, y^\beta [x, y]^\delta, [x, y]^\mu \rangle [F', F],$$

where $\alpha, \beta, \gamma, \delta, \mu \in \mathbb{Z}$. Obviously $p \mid \alpha$ and $p \mid \beta$; if $p \nmid \mu$, then G is abelian. If either $p \nmid \gamma$ or $p \nmid \delta$, then either $\langle x \rangle [G', G]$ or $\langle y \rangle [G', G]$ is normal in $G/[G', G]$ and $G/[G', G]$ is metacyclic; so G is metacyclic.

Finally if $p \mid \gamma, p \mid \mu, p \mid \delta$, then $R \leq F^p [F', F]$, and, if $p = 2$, $R \leq \langle x^2, y^2 \rangle (F')^2 [F', F]$. ■

THEOREM 4.2. Let G be a 2-generator finite p -group.

Then $G \in \mathcal{A}\mathcal{N}_2$ if and only if G is metacyclic and

$$G = \langle a, b \mid |a| = p^\alpha, a^{p^\beta} = b^{p^\gamma}, [a, b] = a^{kp^\delta} \rangle,$$

where $(k, p) = 1, \beta \leq \gamma, \alpha \leq \beta + \delta$ and $\beta \leq \delta$.

PROOF. Such a group is in $\mathcal{A}\mathcal{N}_2$ (see Theorem 3.1).

Now assume that $G = \langle a, b \rangle$ is a p -group in $\mathcal{A}\mathcal{N}_2$.

If G is metacyclic, then G has the required structure (see Theorem 3.1).

Assume for a contradiction that G is non-metacyclic, and let $1 \rightarrow R \rightarrow F \rightarrow G \rightarrow 1$ be a free presentation of G . Thus we have $R \leq F^p [F', F]$ by Lemma 4.1; hence if $p \neq 2$, $[R, F] = [F^p, F][F', F, F] = (F')^p [F', F, F]$ and $[F', F] \leq [R, F]$, contradicting 1.1.

Now let $p = 2$; then for some free generators x, y of F we have $R \leq \langle x^2, y^2 \rangle (F')^2 [F', F]$ and $[R, F] \leq \langle [x^2, y^2], [y^2, x] \rangle (\gamma_3 F)^2 \gamma_4 F = \langle [x, y]^2 [x, y, x], [x, y]^2 [x, y, y] \rangle (\gamma_3 F)^2 \gamma_4 F$. From $[x, y, x] \in [R, F]$ it follows $[x, y, x] = [x, y]^{2b} \cdot [x, y, x]^b [x, y]^{2k} [x, y, x]^k a$, with $a \in (\gamma_3 F)^2 \gamma_4 F$ and $2b + 2k = 0, k \equiv 0(2), b \equiv 1(2)$, contradiction. ■

A finite group is in $\mathcal{A}\mathcal{N}_2$ if and only if its Sylow subgroups are. Furthermore the following result shows that there are no infinite 2-generator non abelian $\mathcal{A}\mathcal{N}_2$ -groups. Hence Theorem 4.2 gives a complete description of 2-generator $\mathcal{A}\mathcal{N}_2$ -groups.

THEOREM 4.3. There are no 2-generator infinite non-abelian $\mathcal{A}\mathcal{N}_2$ -groups.

PROOF. Assume for a contradiction that G is an infinite 2-generator non-abelian $\mathcal{A}\mathcal{N}_2$ -group. Then G/G' is again infinite and 2-generator; we may write $G/G' = \langle xG' \rangle \times \langle yG' \rangle$, with yG' of infinite order. Then $G = \langle x, y \rangle$ and $\langle x \rangle \cap \langle y \rangle = 1$. By 2.4 we can assume $|\langle x, y \rangle| = p$. Thus $x^p, y^p \in \zeta G$ and we can assume that x is a p -element.

If $\langle x \rangle \cap G' = 1$, then the group $\langle x, y \rangle / \langle x^p, y^p \rangle$ is isomorphic to D_4 if $p = 2$ and is not metacyclic if $p \neq 2$, in any case it is not in $\mathcal{A}\mathcal{N}_2$.

If $\langle x \rangle \cap G' = \langle x^{p^a} \rangle \neq 1$, then $|x| = p^{a+1}$ and the group $\langle x, y \rangle / \langle y^{p^{a+1}} \rangle$ is not in $\mathcal{A}\mathcal{N}_2$ by Theorem 4.2. ■

By contrast

THEOREM 4.4. Every infinite 2-generator nil-2 group is residually $A\mathcal{N}_2$.

PROOF. Let G be a 2-generator infinite nil-2 group. Then G/G' is 2-generator and infinite; we can assume that $G/G' = \langle aG' \rangle \times \langle bG' \rangle$, with bG' of infinite order. Thus $G = \langle a, b \rangle$ and $\langle a \rangle \cap \langle b \rangle = 1$. Let T be the torsion subgroup of G , then T is finite and we can assume T to be a p -group (p a prime).

Assume first that $|[a, b]| = p^k$. Then $a^{p^k}, b^{p^k} \in \zeta G$. If a is torsion-free, then, for any $\gamma \geq k$, with $N_\gamma = \langle a^{p^\gamma} [b, a], b^{p^\gamma} [b, a] \rangle$, we have $G/N_\gamma \in A\mathcal{N}_2$; obviously $\cap N_\gamma = 1$ and G is residually $A\mathcal{N}_2$.

If $|a| = p^b$, then $b \geq k$; for any $\gamma \geq b + 1$, with $N_\gamma = \langle b^{p^\gamma} [b, a] \rangle$, we have $G/N_\gamma = \langle b, ab | b^{p^{k+\gamma}} = 1, b^{p^b} = (ab)^{p^b}, [b, ab] = b^{p^\gamma} \rangle$, $b \leq \gamma$ and $G/N_\gamma \in A\mathcal{N}_2$ by Theorem 4.2; moreover $\cap N_\gamma = 1$ as required.

Now let $[a, b]$ be torsion-free; then, with $M_k = \langle [a, b]^{p^k} \rangle$, we have $\cap M_k = 1$ and G/M_k residually $A\mathcal{N}_2$ by the previous case, thus G is residually $A\mathcal{N}_2$ as required. ■

5. INFINITE 3-GENERATOR $A\mathcal{N}_2$ -GROUPS

By 2.1 the direct product of an infinite cyclic group with a finite 2-generator $A\mathcal{N}_2$ -group is in $A\mathcal{N}_2$. The aim of this section is to prove that there are no other infinite 3-generator $A\mathcal{N}_2$ -groups.

THEOREM 5.1. A non-abelian infinite 3-generator group is in $A\mathcal{N}_2$ if and only if it is isomorphic to $C_\infty \times H$, with $H \in A\mathcal{N}_2$, H finite.

PROOF. As remarked above, such groups are in $A\mathcal{N}_2$.

Conversely, assume that $G \in A\mathcal{N}_2$ is a 3-generator infinite group, neither 2-generator nor abelian. Then G/G' is again a 3-generator infinite group and we can write $G/G' = \langle aG' \rangle \times \langle bG' \rangle \times \langle cG' \rangle$, with cG' of infinite order. Then $\langle a, b \rangle \cap \langle c \rangle = 1$. We prove that $G/\langle [a, b] \rangle$ is abelian, and from that it follows $[a, c] = [a, b^\alpha]$, $[b, c] = [a^\beta, b]$ for some $\alpha, \beta \in \mathbb{Z}$ and $[a, ca^\beta b^{-1}] = [b, ca^\beta b^{-\alpha}] = 1$, so $G = \langle a, b \rangle \times \langle cab^{-\alpha} \rangle$, as required.

$G/\langle [a, b] \rangle$ is in $A\mathcal{N}_2$, so we may assume $[a, b] = 1$, $\langle a \rangle \cap \langle b \rangle = 1$ and prove that G is abelian.

It suffices to show that (*) if $[a, c] = 1$, then $[b, c] = 1$.

In fact then $G/\langle [a, c] \rangle$ is abelian, so $[b, c] = [a^\alpha, c]$ for some $\alpha \in \mathbb{Z}$, hence $[ba^{-\alpha}, c] = 1$ and by (*) $[a, c] = 1$.

Then assume $[a, c] = [a, b] = 1$. Since $G/\langle a \rangle$ is a 2-generator infinite $A\mathcal{N}_2$ -group, it follows that $G/\langle a \rangle$ is abelian, by Theorem 4.3, hence $[b, c] = a^\gamma$ for some $\gamma \in \mathbb{Z}$. Then $G = (\langle a \rangle \times \langle b \rangle) \rtimes \langle c \rangle$ and $G = \langle a, b, c | [a, b] = [a, c] = 1, [b, c] = a^\gamma, a^\alpha = 1, b^\beta = 1 \rangle$ for some $\alpha, \beta \in \mathbb{N}_0$ (α or $\beta = 0$ if a or b torsion-free).

Let $1 \rightarrow N \rightarrow F \rightarrow G \rightarrow 1$ be a free presentation of G with $F = \langle x, y, z \rangle$, $N = \langle x^\alpha, y^\beta, [z, y]x^\gamma, [x, y], [x, z] \rangle^F$. Then $N = \langle x^\alpha, y^\beta, [z, y]x^\gamma, [x, y], [x, z], [y, z]^\beta \rangle$.

$\cdot [F', F]$ and $[N, F] = \langle [x, y]^\alpha, [x, z]^\alpha, [y, x]^\beta, [y, z]^\beta, [z, y, z][x, z]^\gamma, [z, y, y][x, y]^\gamma, [x, z, x], [x, z, y], [x, z, z], [x, y, x], [x, y, y], [x, y, z], [z, y, x] \rangle [F', F, F]$.

From $[z, y, z] \in [N, F]$ it follows easily that $[z, y, z] = ([z, y, z][x, z]^\gamma)^\mu ([x, z]^\alpha)^\nu$, and from that $\mu = 1, \gamma = \alpha\nu$, so α divides $\gamma, [b, c] = 1$ and G is abelian, as required. ■

6. HIGHER NILPOTENT EXTENSIONS

Let G be an absolutely nilpotent of class k and H a group possessing a normal subgroup N contained in the n -th term $\zeta_n H$ of the upper central series such that $H/N \cong G$, then $H/[H, N]$ is a central extension by G so has class k , and thus H has class at most $k + n - 1$, one less than is granted by general dispensation. In section 4 we showed that if G is a metacyclic p -group of class k and n any integer ≥ 2 , then there exists an extension $1 \rightarrow N \rightarrow H \rightarrow G \rightarrow 1$, with $N \leq \zeta_n H$ and H of class $s \geq n + k - 1$.

We believe that for most constellations of k and n , every nilpotent group G of class k has an « n -th central extension» $1 \rightarrow N \rightarrow H \rightarrow G \rightarrow 1, N \leq \zeta_n H, n > 1$, such that H is of class at least $k + n - 1$.

We have been able to prove this only for class 2 finite p -groups and for 2-generator p -groups of class ≤ 3 . We start with the following remark:

THEOREM 6.1. Let G be a finite p -group of class $k \geq 2$. Then for any $n \geq k$ there exists a group H of class $\geq n + 1$ with a normal subgroup M such that $M \leq \zeta_n H$ and $H/M \cong G$.

PROOF. Let $1 \rightarrow R \rightarrow F \rightarrow G \rightarrow 1$ be a free presentation of G , with $d(F) = d(G)$, where $d(G)$ is the minimum number of generators of the group G . Then $R \leq F' F^p$, so $[R, F] \leq [F' F^p, F] = [F', F][F^p, F] \leq (F')^p \gamma_3 F$ and, by induction on i , for every $i \geq 1, [R, {}_i F] \leq (F')^p \gamma_{i+2} F$. Therefore, with $N = [R, {}_n F]$, we get $R/N \leq \leq \gamma_n(F/N), (F/N)/(R/N) \cong G$, and the group F/N has class $\geq n + 1$ since $\gamma_{n+1} F \not\leq (F')^p \gamma_{i+2} F$. ■

THEOREM 6.2. Let G be a 2-generator p -group of class ≤ 3 . Then for every integer $n > 1$ there exists a group H of class $n + 2$ with a normal subgroup M such that $M \leq \leq \gamma_n H$ and $H/M \cong G$.

PROOF. If G is metacyclic, the result follows from Theorem 3.1. Otherwise, by Lemma 4.1, there exists a free presentation of $G 1 \rightarrow R \rightarrow F \rightarrow G \rightarrow 1, F = \langle x, y \rangle$ such that $R \leq F' F^p$ and $R \leq \langle x^2, y^2 \rangle (F')^2 [F', F]$ if $p = 2$.

Obviously $\gamma_{n+3} F \leq [R, {}_{n-1} F]$, since $\gamma_4 F \leq R$. We show that, for every $n \geq 1,$

$$(6.1) \quad \gamma_{n+2} F \not\leq [R, {}_n F].$$

From this the result follows. In fact, if $\gamma_{n+3} F \leq [R, {}_n F]$, then the group $H = F/[R, {}_{n-1} F]$ has class $n + 2$ and, with $M = R/[R, {}_{n-1} F]$, we have $M \leq \gamma_{n-1} H \leq \leq \gamma_n H$ and $H/M \cong F/R \cong G$. If $\gamma_{n+3} F \leq [R, {}_n F]$, then the group $H = F/[R, {}_n F]$ has class $n + 2$ and, with $M = R/[R, {}_n F]$ we have $M \leq \gamma_n H$ and $H/M \cong G$.

To establish (6.1) assume first $p \neq 2$. Then $[R, F] \leq [F^p[F', F], F] = [F^p, F]\gamma_4 F = (F')^p \gamma_4 F$ and, for every $n \geq 1$, $[R, {}_n F] \leq (\gamma_{n+1} F)^p \gamma_{n+3} F$. Then $\gamma_{n+2} F \cap [R, {}_n F] \leq \gamma_{n+2} F \cap (\gamma_{n+1} F)^p \gamma_{n+3} F = \gamma_{n+3} F(\gamma_{n+2} F \cap (\gamma_{n+1} F)^p) = \gamma_{n+3} F(\gamma_{n+2} F)^p$. Therefore $\gamma_{n+2} F \not\leq [R, {}_n F]$, since $\gamma_{n+2} F \not\leq \gamma_{n+3} F(\gamma_{n+2} F)^p$.

Assume now $p = 2$. Then $[R, F] \leq \langle [x^2, y], [y^2, x] \rangle^F (\gamma_3 F)^2 \gamma_4 F = \langle [x, y]^2 \cdot [x, y, x], [x, y]^2 [x, y, y] \rangle (\gamma_3 F)^2 \gamma_4 F$, and, by induction on i , it is easy to show that, for every $n \geq 1$, $[R, {}_n F] \leq (\gamma_{n+2} F)^2 \gamma_{n+3} F F'' < d_{i,n} d_{i,n-1}^2, d_{j,n} d_{j-1,n-1}^2 \mid 0 \leq i \leq n-1, 1 \leq j \leq n$.

Assume by contradiction $\gamma_{n+2} F \leq [R, {}_n F]$ for some $n \geq 1$. Then $d_{0,n} \in [N, {}_n F]$ and

$$(6.2) \quad d_{0,n} = \prod_{i=0}^{n-1} d_{i,n}^{b_i} d_{i,n-1}^{2b_i} \prod_{j=1}^n d_{j,n}^{k_j} d_{j-1,n-1}^{2k_j} a$$

for some integers b_i, k_j and $a \in F'' (\gamma_{n+2} F)^2 \gamma_{n+3} F$.

The elements $d_{i,n}, d_{j,n-1}, i \in \{0, \dots, n-1\}, j \in \{0, \dots, n\}$, are free generators of $\gamma_{n+1} F \bmod \gamma_{n+3} F F''$, thus from (6.2) it follows

$$\begin{aligned} 1 &\equiv b_0(2), & 0 &\equiv k_n(2), & b_i + k_i &\equiv 0(2), & \text{for } 1 \leq i \leq n-1, \\ 2b_i + 2k_{i+1} &\equiv 0(2), & & & & & \text{for } 0 \leq i \leq n-1, \end{aligned}$$

and that is impossible. ■

REFERENCES

- [1] F. R. BEYL, *Abelian groups with a vanishing homology group*. J. Pure Appl. Algebra, 7, 1976, 175-193.
- [2] F. R. BEYL, *Isoclinisms of Group Extensions and the Schur Multiplier*. Groups-St. Andrews 1981, London Math. Soc. Lecture Note Ser., vol. 71, Cambridge 1982, 169-185.
- [3] F. R. BEYL, *Commutator properties of extension groups*. C.R. Math. Rep. Acad. Sci. Canada, 2, 1986, 27-30.
- [4] F. R. BEYL - J. TAPPE, *Group Extensions, Representations, and the Schur Multiplier*. Lecture Notes in Mathematics, vol. 958, Springer-Verlag, 1982.
- [5] L. EVENS, *Terminal p -groups*. Illinois J. Math., 12, 1968, 682-699.
- [6] B. H. NEUMANN, *On amalgams of periodic groups*. Proc. Roy. Soc. London, Ser. A 255, 1960, 477-489.
- [7] I. B. S. PASSI - L. R. VERMANI, *The inflation homomorphism*. J. London Math. Soc., II Ser., 6, 1972, 129-136.
- [8] U. STAMMBACH, *Homology in Group Theory*. Lecture Notes in Mathematics, vol. 359, Springer-Verlag, 1973.
- [9] K. VARADARAJAN, *Groups for which Moore spaces $M(\pi, 1)$ exist*. Ann. of Math., 84 (2), 1966, 368-371.

P. Longobardi, M. Maj:

Dipartimento di Matematica e Applicazioni «R. Caccioppoli»
Università degli Studi di Napoli «Federico II»
Complesso Monte S. Angelo
Via Cintia - 80126 NAPOLI

T. MacHenry:

York University
Downsview (12), Ontario (Canada)

J. Wiegold:

School of Mathematics
University of Wales - College of Cardiff
CARDIFF CF2 4AG (Gran Bretagna)