

LIMITS AND COLIMITS OF QUANTALOID-ENRICHED CATEGORIES AND THEIR DISTRIBUTORS

LILI SHEN AND WALTER THOLEN

Dedicated to the memory of Reinhard Börger

ABSTRACT. It is shown that, for a small quantaloid \mathcal{Q} , the category of small \mathcal{Q} -categories and \mathcal{Q} -functors is total and cototal, and so is the category of \mathcal{Q} -distributors and \mathcal{Q} -Chu transforms.

1. INTRODUCTION

The importance of (small) categories enriched in a (unital) quantale rather than in an arbitrary monoidal category was discovered by Lawvere [18] who enabled us to look at individual mathematical objects, such as metric spaces, as small categories. Through the study of lax algebras [15], quantale-enriched categories have become the backbone of a larger array of objects that may be viewed as individual generalized categories. Prior to this development, Walters [31] had extended Lawvere’s viewpoint in a different manner, replacing the quantale at work by a *quantaloid* (a term proposed later by Rosenthal [22]), thus by a bicategory with the particular property that its hom-objects are given by complete lattices such that composition from either side preserves suprema; quantales are thus simply one-object quantaloids.

Based on the theory of quantaloid-enriched categories developed by Stubbe [25, 26], recent works [16, 21, 27, 28] have considered in particular the case when the quantaloid in question arises from a given quantaloid by a “diagonal construction” whose roots go far beyond its use in this paper; see [12, 13]. Specifically for the one-object quantaloids (i.e., quantales) whose enriched categories give (pre)ordered sets and (generalized) metric spaces, the corresponding small quantaloids of diagonals lead to truly partial structures, in the sense that the full structure is available only on a subset of the ambient underlying set of objects.

In the first instance then, this paper aims at exploring the categorical properties of the category $\mathcal{Q}\text{-Cat}$ of small \mathcal{Q} -enriched categories and their \mathcal{Q} -functors for a small quantaloid \mathcal{Q} . By showing that $\mathcal{Q}\text{-Cat}$ is topological [2] over the comma category $\mathbf{Set}/\text{ob } \mathcal{Q}$ (Proposition 2.2) one easily describes small limits and colimits in this category, and beyond. In fact, one concludes that categories of this type are total [24] and cototal, hence possess even those limits and colimits of large diagrams whose existence is not made impossible by the size of the small hom-sets of $\mathcal{Q}\text{-Cat}$ (see [9]).

Our greater interest, however, is in the category $\mathcal{Q}\text{-Chu}$ whose objects are often called \mathcal{Q} -Chu spaces, the prototypes of which go back to [3, 20] and many others (see [4, 10]). Its objects are \mathcal{Q} -distributors of \mathcal{Q} -categories (also called \mathcal{Q} -(bi)modules or \mathcal{Q} -profunctors), hence they are compatible \mathcal{Q} -relations (or \mathcal{Q} -matrices) that have been investigated intensively ever since Bénabou [5] introduced them (see [6, 7]). While when taken as the morphisms of the category whose objects are \mathcal{Q} -categories, they make for a in many ways poorly performing category (as already the case $\mathcal{Q} = \mathbf{2}$ shows), when taken as objects of $\mathcal{Q}\text{-Chu}$ with morphisms given by so-called \mathcal{Q} -Chu transforms, i.e., by pairs of \mathcal{Q} -functors that behave like adjoint operators, we obtain a category that in terms of the existence of limits and colimits behaves as strongly as $\mathcal{Q}\text{-Cat}$ itself. In analogy to the property shown in [11] in a different categorical context, we first prove that the domain functor $\mathcal{Q}\text{-Chu} \rightarrow \mathcal{Q}\text{-Cat}$ allows for initial liftings [2] of structured cones over small diagrams

2010 *Mathematics Subject Classification.* 18D20, 18A30.

Partial financial assistance by the Natural Sciences and Engineering Research Council of Canada is gratefully acknowledged.

(Theorem 3.4), which then allows for an explicit description of all limits and colimits in $\mathcal{Q}\text{-Chu}$ over small diagrams. But although the domain functor fails to be topological, just as for $\mathcal{Q}\text{-Cat}$ we are able to show totality (and, consequently, cototality) of $\mathcal{Q}\text{-Chu}$. A key ingredient for this result is the existence proof of a generating set in $\mathcal{Q}\text{-Chu}$, and therefore also of a cogenerating set (Theorem 3.8).

2. LIMITS AND COLIMITS IN $\mathcal{Q}\text{-Cat}$

Throughout, let \mathcal{Q} be a small *quantaloid*, i.e., a small category enriched in the category **Sup** of complete lattices and sup-preserving maps. A small \mathcal{Q} -category is given by a set X (its set of objects) and a lax functor

$$a : X \longrightarrow \mathcal{Q},$$

where the set X is regarded as a quantaloid carrying the chaotic structure, so that for all $x, y \in X$ there is precisely one arrow $x \longrightarrow y$, called (x, y) . Explicitly then, the \mathcal{Q} -category structure on X is given by

- a family of objects $|x|_X := ax$ in \mathcal{Q} ($x \in X$),
- a family of morphisms $a(x, y) : |x| \longrightarrow |y|$ in \mathcal{Q} ($x, y \in X$), subject to

$$1_{|x|} \leq a(x, x) \quad \text{and} \quad a(y, z) \circ a(x, y) \leq a(x, z)$$

($x, y, z \in X$). When one calls $|x| = |x|_X$ the *extent* (or *type*) of $x \in X$, a \mathcal{Q} -functor $f : (X, a) \longrightarrow (Y, b)$ of \mathcal{Q} -categories (X, a) , (Y, b) is an extent-preserving map $f : X \longrightarrow Y$ such that there is a lax natural transformation $a \longrightarrow bf$ given by identity morphisms in \mathcal{Q} ; explicitly,

$$|x|_X = |f(x)|_Y \quad \text{and} \quad a(x, y) \leq b(f(x), f(y))$$

for all $x, y \in X$. Denoting the resulting ordinary category by $\mathcal{Q}\text{-Cat}$, we have a forgetful functor

$$\begin{array}{ccc} \mathcal{Q}\text{-Cat} & \xrightarrow{\text{ob}} & \mathbf{Set}/\text{ob } \mathcal{Q} \\ \begin{array}{ccc} X & \xrightarrow{f} & Y \\ & \searrow a & \swarrow b \\ & \mathcal{Q} & \end{array} & \mapsto & \begin{array}{ccc} X & \xrightarrow{f} & Y \\ & \searrow |\cdot| & \swarrow |\cdot| \\ & \text{ob } \mathcal{Q} & \end{array} \end{array}$$

Example 2.1. (1) If \mathcal{Q} is a *quantale*, i.e., a one-object quantaloid, then the extent functions of \mathcal{Q} -categories are trivial and $\mathcal{Q}\text{-Cat}$ assumes its classical meaning (as in [17], where \mathcal{Q} is considered as a monoidal (closed) category). Prominent examples are $\mathcal{Q} = \mathbf{2} = \{0 < 1\}$ and $\mathcal{Q} = ([0, \infty], \geq, +)$, where then $\mathcal{Q}\text{-Cat}$ is the category **Ord** (sets carrying a reflexive and transitive relation, called *order* here but commonly known as preorder, with monotone maps) and, respectively, the category **Met** (sets X carrying a distance function $a : X \times X \longrightarrow [0, \infty]$ required to satisfy only $a(x, x) = 0$ and $a(x, z) \leq a(x, y) + a(y, z)$ but, in accordance with the terminology introduced by Lawvere [18] and used in [15], nevertheless called *metric* here, with non-expanding maps $f : X \longrightarrow Y$, so that $b(f(x), f(y)) \leq a(x, y)$ for all $x, y \in X$).

- (2) (Stubbe [27]) Every quantaloid \mathcal{Q} gives rise to a new quantaloid $\mathbf{D}\mathcal{Q}$ whose objects are the morphisms of \mathcal{Q} , and for morphisms u, v in \mathcal{Q} , a morphism $(u, d, v) : u \rightsquigarrow v$ in $\mathbf{D}\mathcal{Q}$, normally written just as d , is a \mathcal{Q} -morphism $d : \text{dom } u \longrightarrow \text{cod } v$ satisfying

$$(d \swarrow u) \circ u = d = v \circ (v \searrow d),$$

also called a *diagonal* from u to v :

$$\begin{array}{ccc} \bullet & \xrightarrow{v \searrow d} & \bullet \\ \downarrow u & \searrow d & \downarrow v \\ \bullet & \xrightarrow{d \swarrow u} & \bullet \end{array}$$

(Here $d \swarrow u, v \searrow d$ denote the *internal homs* of \mathcal{Q} , determined by

$$z \leq d \swarrow u \iff z \circ u \leq d, \quad t \leq v \searrow d \iff v \circ t \leq d$$

for all $z : \text{cod } u \rightarrow \text{cod } d, t : \text{dom } d \rightarrow \text{dom } v$.) With the composition of $d : u \rightsquigarrow v$ with $e : v \rightsquigarrow w$ in \mathbf{DQ} defined by

$$e \diamond d = (e \swarrow v) \circ d = e \circ (v \searrow d),$$

and with identity morphisms $u : u \rightsquigarrow u$, \mathbf{DQ} becomes a quantaloid whose local order is inherited from \mathcal{Q} . In fact, there is a full embedding

$$\mathcal{Q} \rightarrow \mathbf{DQ}, \quad (u : t \rightarrow s) \mapsto (u : 1_t \rightsquigarrow 1_s)$$

of quantaloids.

We remark that the construction of \mathbf{D} works for ordinary categories; indeed it is part of the proper factorization monad on \mathbf{CAT} [13].

- (2a) For $\mathcal{Q} = \mathbf{2}$, the quantaloid \mathbf{DQ} has object set $\{0, 1\}$. There are exactly two \mathbf{DQ} -arrows $1 \rightsquigarrow 1$, given by $0, 1$, and 0 is the only arrow in every other hom-set of \mathbf{DQ} ; composition is given by infimum. A \mathbf{DQ} -category is given by a set X , a distinguished subset $A \subseteq X$ (those elements of X with extent 1) and a (pre)order on A . Hence, a \mathbf{DQ} -category structure is a (truly!) *partial order*, and with those maps that may be monotonely restricted to the distinguished subsets as morphisms. We write

$$\mathbf{ParOrd} = \mathbf{DQ-Cat}$$

for the resulting category, which contains \mathbf{Ord} as a full coreflective subcategory.

- (2b) For $\mathcal{Q} = ([0, \infty], \geq, +)$, the hom-sets of \mathbf{DQ} are easily described by

$$\mathbf{DQ}(u, v) = \{s \in [0, \infty] \mid u, v \leq s\},$$

with composition given by $t \diamond s = s - v + t$ (for $t : v \rightsquigarrow w$). A \mathbf{DQ} -category structure on a set X consists of functions $|\cdot| : X \rightarrow [0, \infty], a : X \times X \rightarrow [0, \infty]$ satisfying

$$|x|, |y| \leq a(x, y), \quad a(x, x) \leq |x|, \quad a(x, z) \leq a(x, y) - |y| + a(y, z)$$

($x, y, z \in X$). Obviously, since necessarily $|x| = a(x, x)$, these conditions simplify to

$$a(x, x) \leq a(x, y), \quad a(x, z) \leq a(x, y) - a(y, y) + a(y, z)$$

($x, y, z \in X$), describing a as a *partial metric* on X (see [16, 19, 21]¹). With non-expanding maps one obtains the category

$$\mathbf{ParMet} = \mathbf{DQ-Cat},$$

which contains \mathbf{Met} as a full coreflective subcategory: the coreflector restricts the partial metric a on X to those elements $x \in X$ with $a(x, x) = 0$.

To see how limits and colimits in the (ordinary) category $\mathbf{Q-Cat}$ are to be formed, it is best to first prove its topologicity over $\mathbf{Set}/\text{ob } \mathcal{Q}$. Recall that, for any functor $U : \mathcal{A} \rightarrow \mathcal{X}$, a *U-structured cone* over a diagram $D : \mathcal{J} \rightarrow \mathcal{A}$ is given by an object $X \in \mathcal{X}$ and a natural transformation $\xi : \Delta X \rightarrow UD$. A *lifting* of (X, ξ) is given by an object A in \mathcal{A} and a cone $\alpha : \Delta A \rightarrow D$ over D with $UA = X, U\alpha = \xi$. Such lifting (A, α) is *U-initial* if, for all cones $\beta : \Delta B \rightarrow D$ over D and morphisms $t : UB \rightarrow UA$ in \mathcal{X} , there is exactly one morphism $h : B \rightarrow A$ in \mathcal{A} with $Uh = t$ and $\alpha \cdot \Delta h = \beta$. We call U *small-topological* [11] if all U -structured cones over small diagrams admit U -initial liftings, and U is *topological* when this condition holds without the size restriction on diagrams. Recall also the following well-known facts:

- Topological functors are necessarily faithful [8], and for faithful functors it suffices to consider discrete cones to guarantee topologicity.
- $U : \mathcal{A} \rightarrow \mathcal{X}$ is topological if, and only if, $U^{\text{op}} : \mathcal{A}^{\text{op}} \rightarrow \mathcal{X}^{\text{op}}$ is topological.

¹Here our terminology naturally extends Lawvere's notion of metric and is synonymous with "generalized partial metric" as used by Pu-Zhang [21] who dropped finiteness ($a(x, y) < \infty$), symmetry ($a(x, y) = a(y, x)$) and separation ($a(x, x) = a(x, y) = a(y, y) \iff x = y$) from the requirement for the notion of "partial metric" as originally introduced by Matthews [19].

- The two properties above generally fail to hold for small-topological functors. However, for any functor U , a U -initial lifting of a U -structured cone that is a limit cone in \mathcal{X} gives also a limit cone in \mathcal{A} .
- Every small-topological functor is a fibration (consider singleton diagrams) and has a fully faithful right adjoint (consider the empty diagram).

Proposition 2.2. *For every (small) quantaloid \mathcal{Q} , the “object functor” $\mathcal{Q}\text{-Cat} \rightarrow \mathbf{Set}/\text{ob } \mathcal{Q}$ is topological.*

Proof. Given a (possibly large) family $f_i : (X, |-|) \rightarrow (Y_i, |-|_i)$ ($i \in I$) of maps over $\text{ob } \mathcal{Q}$, where every Y_i carries a \mathcal{Q} -category structure b_i with extent function $|-|_i$, we must find a \mathcal{Q} -category structure a on X with extent function $|-|$ such that (1) every $f_i : (X, a) \rightarrow (Y, b)$ is a \mathcal{Q} -functor, and (2) for every \mathcal{Q} -category (Z, c) , any extent preserving map $g : Z \rightarrow X$ becomes a \mathcal{Q} -functor $(Z, c) \rightarrow (X, a)$ whenever all maps $f_i g$ are \mathcal{Q} -functors $(Z, c) \rightarrow (Y_i, b_i)$ ($i \in I$). But this is easy: simply define

$$a(x, y) := \bigwedge_{i \in I} b_i(f_i(x), f_i(y))$$

for all $x, y \in X$. Hence, a is the ob-initial structure on X with respect to the structured sink $(f_i : X \rightarrow (Y_i, b_i))_{i \in I}$. \square

Corollary 2.3. *$\mathcal{Q}\text{-Cat}$ is complete and cocomplete, and the object functor has both a fully faithful left adjoint and a fully faithful right adjoint.*

Remark 2.4. (1) The set of objects of the product (X, a) of a small family of \mathcal{Q} -categories (X_i, a_i) ($i \in I$) is given by the fibred product of $(X_i, |-|_i)$ ($i \in I$), i.e.,

$$X = \{((x_i)_{i \in I}, q) \mid q \in \text{ob } \mathcal{Q}, \forall i \in I (x_i \in X_i, |x_i| = q)\},$$

and (when writing $(x_i)_{i \in I}$ instead of $((x_i)_{i \in I}, q)$ and putting $|(x_i)_{i \in I}| = q$) we have

$$a((x_i)_{i \in I}, (y_i)_{i \in I}) = \bigwedge_{i \in I} a_i(x_i, y_i) : |(x_i)_{i \in I}| \rightarrow |(y_i)_{i \in I}|$$

for its hom-arrows. In particular, $(\text{ob } \mathcal{Q}, \top)$ with

$$\top(q, r) = \top : q \rightarrow r$$

the top element in $\mathcal{Q}(q, r)$ (for all $q, r \in \text{ob } \mathcal{Q}$), is the terminal object in $\mathcal{Q}\text{-Cat}$.

- (2) The coproduct (X, a) of \mathcal{Q} -categories (X_i, a_i) ($i \in I$) is simply formed by the coproduct in \mathbf{Set} , with all structure to be obtained by restriction:

$$X = \coprod_{i \in I} X_i, \quad |x|_X = |x|_{X_i} \text{ if } x \in X_i, \quad a(x, y) = \begin{cases} a_i(x, y) & \text{if } x, y \in X_i, \\ \perp : |x| \rightarrow |y| & \text{else.} \end{cases}$$

In particular, \emptyset with its unique \mathcal{Q} -category structure is an initial object in $\mathcal{Q}\text{-Cat}$.

- (3) The equalizer of \mathcal{Q} -functors $f, g : (X, a) \rightarrow (Y, b)$ is formed as in \mathbf{Set} , by restriction of the structure of (X, a) . The object set of their coequalizer (Z, c) in $\mathcal{Q}\text{-Cat}$ is also formed as in \mathbf{Set} , so that $Z = Y / \sim$, with \sim the least equivalence relation on Y with $f(x) \sim g(x)$, $x \in X$. With $\pi : Y \rightarrow Z$ the projection, necessarily $|\pi(y)|_Z = |y|_Y$, and $c(\pi(y), \pi(y'))$ is the join of all

$$b(y_n, y'_n) \circ b(y_{n-1}, y'_{n-1}) \circ \cdots \circ b(y_2, y'_2) \circ b(y_1, y'_1),$$

where $|y| = |y_1|, |y'_1| = |y_2|, \dots, |y'_{n-1}| = |y_n|, |y'_n| = |y'|$ ($y_i, y'_i \in Y, n \geq 1$).

- (4) The fully faithful left adjoint of $\mathcal{Q}\text{-Cat} \rightarrow \mathbf{Set}/\text{ob } \mathcal{Q}$ provides a set $(X, |-|)$ over $\text{ob } \mathcal{Q}$ with the discrete \mathcal{Q} -structure, given by

$$a(x, y) = \begin{cases} 1_{|x|} & \text{if } x = y, \\ \perp : |x| \rightarrow |y| & \text{else;} \end{cases}$$

while the fully faithful right adjoint always takes $\top : |x| \rightarrow |y|$ as the hom-arrow, i.e., it chooses the indiscrete \mathcal{Q} -structure.

Example 2.5. The product of partial metric spaces (X_i, a_i) ($i \in I$) provides its carrier set

$$X = \{((x_i)_{i \in I}, s) \mid s \in [0, \infty], \forall i \in I (x_i \in X_i, |x_i| = s)\}$$

with the “sup metric”:

$$a((x_i)_{i \in I}, (y_i)_{i \in I}) = \sup_{i \in I} a_i(x_i, y_i).$$

$[0, \infty]$ is terminal in **ParMet** when provided with the chaotic metric that makes all distances 0, and it is a generator when provided with the discrete metric d :

$$d(s, t) = \begin{cases} 0 & \text{if } s = t, \\ \infty & \text{else.} \end{cases}$$

Beyond small limits and colimits, **Q-Cat** actually has all large-indexed limits and colimits that one can reasonably expect to exist. More precisely, recall that an ordinary category \mathcal{C} with small hom-sets is (see [9])

- *hypercomplete* if a diagram $D : \mathcal{J} \rightarrow \mathcal{C}$ has a limit in \mathcal{C} whenever the limit of $\mathcal{C}(A, D-)$ exists in **Set** for all $A \in \text{ob } \mathcal{C}$; equivalently: whenever, for every $A \in \text{ob } \mathcal{C}$, the cones $\Delta A \rightarrow D$ in \mathcal{C} may be labeled by a set;
- *totally cocomplete* if a diagram $D : \mathcal{J} \rightarrow \mathcal{C}$ has a colimit in \mathcal{C} whenever the colimit of $\mathcal{C}(A, D-)$ exists in **Set** for all $A \in \text{ob } \mathcal{C}$; equivalently: whenever, for every $A \in \text{ob } \mathcal{C}$, the connected components of $(A \downarrow D)$ may be labelled by a set.

The dual notions are *hypercocomplete* and *totally complete*. It is well known (see [9]) that

- \mathcal{C} is totally cocomplete if, and only if, \mathcal{C} is *total*, i.e., if the Yoneda embedding $\mathcal{C} \rightarrow \mathbf{Set}^{\mathcal{C}^{\text{op}}}$ has a left adjoint;
- total cocompleteness implies hypercompleteness but not vice versa (with Adámek’s monadic category over graphs [1] providing a counterexample);
- for a *solid* (=semi-topological [29]) functor $\mathcal{A} \rightarrow \mathcal{X}$, if \mathcal{X} is hypercomplete or totally complete, \mathcal{A} has the corresponding property [30];
- in particular, every topological functor, every monadic functor over **Set**, and every full reflective embedding is solid.

It is also useful for us to recall [9, Corollary 3.5]:

Proposition 2.6. *A cocomplete and cowellpowered category with small hom-sets and a generating set of objects is total.*

Since **Q-Cat** is topological over **Set/ob Q** which, as a complete, cocomplete, wellpowered and cowellpowered category with a generating and a cogenerating set, is totally complete and totally cocomplete, we conclude:

Theorem 2.7. *Q-Cat is totally complete and totally cocomplete and, in particular, hypercocomplete and hypercomplete.*

Remark 2.8. Of course, we may also apply Proposition 2.6 directly to obtain Theorem 2.7 since the left adjoint of **Q-Cat** \rightarrow **Set/ob Q** sends a generating set of **Set/ob Q** to a generating set of **Q-Cat**, and the right adjoint has the dual property. Explicitly then, denoting for every $s \in \text{ob } \mathcal{Q}$ by $\{s\}$ the discrete \mathcal{Q} -category whose only object has extent s , we obtain the generating set $\{\{s\} \mid s \in \text{ob } \mathcal{Q}\}$ for **Q-Cat**. Similarly, providing the disjoint unions $D_s = \{s\} + \text{ob } \mathcal{Q}$ ($s \in \text{ob } \mathcal{Q}$) with the identical extent functions and the indiscrete \mathcal{Q} -category structures, one obtains a cogenerating set in **Q-Cat**.

3. LIMITS AND COLIMITS IN **Q-Chu**

For \mathcal{Q} -categories $X = (X, a)$, $Y = (Y, b)$, a \mathcal{Q} -distributor [5] $\varphi : X \dashv\!\!\dashv Y$ (also called \mathcal{Q} -(bi)module [18], \mathcal{Q} -profunctor) is a family of arrows $\varphi(x, y) : |x| \rightarrow |y|$ ($x \in X, y \in Y$) in \mathcal{Q} such that

$$b(y, y') \circ \varphi(x, y) \circ a(x', x) \leq \varphi(x', y')$$

for all $x, x' \in X, y, y' \in Y$. Its composite with $\psi : Y \dashrightarrow Z$ is given by

$$(\psi \circ \varphi)(x, z) = \bigvee_{y \in Y} \psi(y, z) \circ \varphi(x, y).$$

Since the structure a of a \mathcal{Q} -category (X, a) is neutral with respect to this composition, we obtain the category

$\mathcal{Q}\text{-Dis}$

of \mathcal{Q} -categories and their \mathcal{Q} -distributors which, with the local pointwise order

$$\varphi \leq \varphi' \iff \forall x, y : \varphi(x, y) \leq \varphi'(x, y),$$

is actually a quantaloid. Every \mathcal{Q} -functor $f : X \rightarrow Y$ gives rise to the \mathcal{Q} -distributors $f_{\natural} : X \dashrightarrow Y$ and $f^{\natural} : Y \dashrightarrow X$ with

$$f_{\natural}(x, y) = b(f(x), y) \quad \text{and} \quad f^{\natural}(y, x) = b(y, f(x))$$

($x \in X, y \in Y$). One has $f_{\natural} \dashv f^{\natural}$ in the 2-category $\mathcal{Q}\text{-Dis}$, and if one lets $\mathcal{Q}\text{-Cat}$ inherit the order of $\mathcal{Q}\text{-Dis}$ via

$$f \leq g \iff f_{\natural} \leq g_{\natural} \iff g^{\natural} \leq f^{\natural} \iff 1_{|x|} \leq b(f(x), g(x)) \quad (x \in X),$$

then one obtains 2-functors

$$(-)_{\natural} : (\mathcal{Q}\text{-Cat})^{\text{co}} \rightarrow \mathcal{Q}\text{-Dis}, \quad (-)^{\natural} : (\mathcal{Q}\text{-Cat})^{\text{op}} \rightarrow \mathcal{Q}\text{-Dis}$$

which map objects identically; here ‘‘op’’ refers to the dualization of 1-cells and ‘‘co’’ to the dualization of 2-cells.

Example 3.1 (See Example 2.1). (1) A **2**-distributor is an *order ideal relation*; that is, a relation $\varphi : X \dashrightarrow Y$ of ordered sets that behaves like a two-sided ideal w.r.t. the order:

$$x' \leq x \ \& \ x \varphi y \ \& \ y \leq y' \implies x' \varphi y'.$$

A $[0, \infty]$ -distributor $\varphi : X \dashrightarrow Y$ introduces a distance function between metric spaces $(X, a), (Y, b)$ that must satisfy

$$\varphi(x', y') \leq a(x', x) + \varphi(x, y) + a(y, y')$$

for all $x, x' \in X, y, y' \in Y$.

- (2) A **D2**-distributor $\varphi : X \dashrightarrow Y$ is given by a **2**-distributor $A \dashrightarrow B$ where $A = \{x \in X \mid x \leq x\}, B = \{y \in Y \mid y \leq y\}$ are the coreflections of X, Y , respectively. Likewise, a $\text{D}[0, \infty]$ -distributor $\varphi : X \dashrightarrow Y$ is given by a distributor of the metric coreflections of the partial metric spaces X and Y .

In our context $\mathcal{Q}\text{-Dis}$ plays only an auxiliary role for us in setting up the category

$\mathcal{Q}\text{-Chu}$

whose objects are \mathcal{Q} -distributors and whose morphisms $(f, g) : \varphi \rightarrow \psi$ are given by \mathcal{Q} -functors $f : (X, a) \rightarrow (Y, b), g : (Z, c) \rightarrow (W, d)$ such that the diagram

$$\begin{array}{ccc} X & \xrightarrow{f_{\natural}} & Y \\ \varphi \circ \downarrow & & \downarrow \psi \\ W & \xrightarrow{g^{\natural}} & Z \end{array} \quad (3.i)$$

commutes in $\mathcal{Q}\text{-Dis}$:

$$\psi(f(x), z) = \varphi(x, g(z)) \quad (3.ii)$$

for all $x \in X, z \in Z$. In particular, with $\varphi = a, \psi = b$ one obtains that the morphisms $(f, g) : 1_{(X, a)} \rightarrow 1_{(Y, b)}$ in $\mathcal{Q}\text{-Chu}$ are precisely the adjunctions $f \dashv g : (Y, b) \rightarrow (X, a)$ in the

2-category $\mathcal{Q}\text{-Cat}$. With the order inherited from $\mathcal{Q}\text{-Cat}$, $\mathcal{Q}\text{-Chu}$ is in fact a 2-category, and one has 2-functors

$$\begin{aligned} \text{dom} : \mathcal{Q}\text{-Chu} &\longrightarrow \mathcal{Q}\text{-Cat}, & (f, g) &\mapsto f, \\ \text{cod} : \mathcal{Q}\text{-Chu} &\longrightarrow (\mathcal{Q}\text{-Cat})^{\text{op}}, & (f, g) &\mapsto g. \end{aligned}$$

In order for us to exhibit properties of $\mathcal{Q}\text{-Chu}$, it is convenient to describe $\mathcal{Q}\text{-Chu}$ transforms, i.e., morphisms in $\mathcal{Q}\text{-Chu}$, alternatively, with the help of *presheaves*, as follows. For every $s \in \text{ob } \mathcal{Q}$, let $\{s\}$ denote the discrete \mathcal{Q} -category whose only object has extent s . For a \mathcal{Q} -category $X = (X, a)$, a \mathcal{Q} -presheaf φ on X of extent $|\varphi| = s$ is a \mathcal{Q} -distributor $\varphi : X \dashrightarrow \{s\}$. Hence, φ is given by a family of \mathcal{Q} -morphisms $\varphi_x : |x| \longrightarrow |\varphi|$ ($x \in X$) with $\varphi_y \circ a(x, y) \leq \varphi_x$ ($x, y \in X$). With

$$[\varphi, \psi] = \bigwedge_{x \in X} \psi_x \swarrow \varphi_x,$$

PX becomes a \mathcal{Q} -category, and one has the *Yoneda \mathcal{Q} -functor*

$$y_X = y : X \longrightarrow \text{PX}, \quad x \mapsto (a(-, x) : X \dashrightarrow \{|x|\}).$$

y is fully faithful, i.e., $[y(x), y(y)] = a(x, y)$ ($x, y \in X$). The point of the formation of PX for us is as follows (see [14, 23]):

Proposition 3.2. *The 2-functor $(-)^{\natural} : (\mathcal{Q}\text{-Cat})^{\text{op}} \longrightarrow \mathcal{Q}\text{-Dis}$ has a left adjoint \mathbb{P} which maps a \mathcal{Q} -distributor $\varphi : X \dashrightarrow Y$ to the \mathcal{Q} -functor*

$$\varphi^* : \text{PY} \longrightarrow \text{PX}, \quad \psi \mapsto \psi \circ \varphi;$$

hence,

$$(\varphi^*(\psi))_x = \bigvee_{y \in Y} \psi_y \circ \varphi(x, y)$$

for all $\psi \in \text{PY}$, $x \in X$. In particular, for a \mathcal{Q} -functor $f : X \longrightarrow Y$ one has

$$f^* := (f_{\natural})^* : \text{PY} \longrightarrow \text{PX}, \quad (f^*(\psi))_x = \psi_{f(x)}.$$

Denoting by $\tilde{\varphi} : Y \longrightarrow \text{PX}$ the transpose of $\varphi : X \dashrightarrow Y$ under the adjunction, determined by $\tilde{\varphi}^{\natural} \circ (y_X)_{\natural} = \varphi$, so that $(\tilde{\varphi}(y))_x = \varphi(x, y)$ for all $x \in X$, $y \in Y$, we can now present $\mathcal{Q}\text{-Chu}$ transforms, as follows:

Corollary 3.3. *A morphism $(f, g) : \varphi \longrightarrow \psi$ in $\mathcal{Q}\text{-Chu}$ (as in (3.i)) may be equivalently presented as a commutative diagram*

$$\begin{array}{ccc} \text{PX} & \xleftarrow{f^*} & \text{PY} \\ \tilde{\varphi} \uparrow & & \uparrow \tilde{\psi} \\ \text{W} & \xleftarrow{g} & \text{Z} \end{array} \quad (3.iii)$$

in $\mathcal{Q}\text{-Cat}$. Condition (3.ii) then reads as

$$(\tilde{\varphi}(g(z)))_x = (\tilde{\psi}(z))_{f(x)} \quad (3.iv)$$

for all $x \in X$, $z \in Z$.

Proof. For all $z \in Z$,

$$f^*(\tilde{\psi}(z)) = \tilde{\psi}(z) \circ f_{\natural} = y_Z(z) \circ \psi \circ f_{\natural} = y_Z(z) \circ g^{\natural} \circ \varphi = y_Y(g(z)) \circ \varphi = \tilde{\varphi}(g(z)).$$

□

Theorem 3.4. *Let $D : \mathcal{J} \longrightarrow \mathcal{Q}\text{-Chu}$ be a diagram such that the colimit $W = \text{colim cod } D$ exists in $\mathcal{Q}\text{-Cat}$. Then any cone $\gamma : \Delta X \longrightarrow \text{dom } D$ in $\mathcal{Q}\text{-Cat}$ has a dom-initial lifting $\Gamma : \Delta \varphi \longrightarrow D$ in $\mathcal{Q}\text{-Chu}$ with $\varphi : X \dashrightarrow W$, $\text{dom } \Gamma = \gamma$. In particular, if γ is a limit cone in $\mathcal{Q}\text{-Cat}$, Γ is a limit cone in $\mathcal{Q}\text{-Chu}$.*

Proof. Considering the functors

$$\mathcal{Q}\text{-Chu} \xrightarrow{\text{dom}} \mathcal{Q}\text{-Cat} \xrightarrow{(-)^\natural} \mathcal{Q}\text{-Dis}, \quad \mathcal{Q}\text{-Chu} \xrightarrow{\text{cod}} (\mathcal{Q}\text{-Cat})^{\text{op}} \xrightarrow{(-)^\natural} \mathcal{Q}\text{-Dis},$$

one has the natural transformation

$$\kappa : (\text{dom}(-))^\natural \dashv \Rightarrow (\text{cod}(-))^\natural, \quad \kappa_\varphi := \varphi \ (\varphi \in \text{ob } \mathcal{Q}\text{-Chu}).$$

By the adjunction of Proposition 3.2, $\kappa D : (\text{dom } D)^\natural \dashv \Rightarrow (\text{cod } D)^\natural$ corresponds to a natural transformation $\widetilde{\kappa D} : \text{cod } D \rightarrow \mathbf{P}(\text{dom } D)^\natural$, and the given cone γ gives a cocone $\gamma^* : \mathbf{P}(\text{dom } D)^\natural \rightarrow \Delta \mathbf{P}X$. Forming the colimit cocone $\delta : \text{cod } D \rightarrow \Delta W$ one now obtains a unique \mathcal{Q} -functor $\widetilde{\varphi} : W \rightarrow \mathbf{P}X$ making

$$\begin{array}{ccc} \Delta \mathbf{P}X & \xleftarrow{\gamma^*} & \mathbf{P}(\text{dom } D)^\natural \\ \uparrow \Delta \widetilde{\varphi} & & \uparrow \widetilde{\kappa D} \\ \Delta W & \xleftarrow{\delta} & \text{cod } D \end{array}$$

commute in $\mathcal{Q}\text{-Cat}$ or, equivalently, making

$$\begin{array}{ccc} \Delta X & \xrightarrow{\gamma^\natural} & (\text{dom } D)^\natural \\ \downarrow \Delta \varphi & & \downarrow \kappa D \\ \Delta W & \xrightarrow{\delta^\natural} & (\text{cod } D)^\natural \end{array}$$

commute in $\mathcal{Q}\text{-Dis}$, with $\varphi : X \dashv \Rightarrow W$ corresponding to $\widetilde{\varphi}$. In other words, we have a cone $\Gamma : \Delta \varphi \rightarrow D$ with $\text{dom } \varphi = X$, $\text{dom } \Gamma = \gamma$, namely $\Gamma = (\gamma, \delta)$.

Given a cone $\Theta : \Delta \psi \rightarrow D$ with $\psi : Y \dashv \Rightarrow Z$ in $\mathcal{Q}\text{-Dis}$ and a \mathcal{Q} -functor $f : Y \rightarrow X$ with $\gamma \cdot \Delta f = \epsilon := \text{dom } \Theta$, the cocone $\vartheta := \text{cod } \Theta : \text{cod } D \rightarrow \Delta Z$ corresponds to a unique \mathcal{Q} -functor $g : W \rightarrow Z$ with $\Delta g \cdot \delta = \vartheta$ by the colimit property. As the diagram

$$\begin{array}{ccccc} & & \Delta \mathbf{P}X & \xleftarrow{\gamma^*} & \mathbf{P}(\text{dom } D)^\natural \\ & \Delta f^* \nearrow & \uparrow \Delta \widetilde{\varphi} & \epsilon^* \nearrow & \uparrow \widetilde{\kappa D} \\ \Delta \mathbf{P}Y & & \Delta W & \xleftarrow{\delta} & \text{cod } D \\ \uparrow \Delta \widetilde{\psi} & \Delta g \nearrow & & \vartheta \nearrow & \\ \Delta Z & & & & \end{array}$$

shows, the colimit property of W also guarantees $f^* \widetilde{\varphi} = \widetilde{\psi} g$ (with $\widetilde{\psi}$ corresponding to ψ) which, by Corollary 3.3, means that $(f, g) : \psi \rightarrow \varphi$ is the only morphism in $\mathcal{Q}\text{-Chu}$ with $\text{dom}(f, g) = f$ and $\Gamma \cdot \Delta(f, g) = \Theta$. \square

Corollary 3.5. *dom : $\mathcal{Q}\text{-Chu} \rightarrow \mathcal{Q}\text{-Cat}$ is small-topological; in particular, dom is a fibration with a fully faithful right adjoint which embeds $\mathcal{Q}\text{-Cat}$ into $\mathcal{Q}\text{-Chu}$ as a full reflective subcategory. cod : $\mathcal{Q}\text{-Chu} \rightarrow (\mathcal{Q}\text{-Cat})^{\text{op}}$ has the dual properties.*

Proof. With the existence of small colimits guaranteed by Corollary 2.3, dom-initial liftings to small dom-structured cones exist by Theorem 3.4. For the assertion on cod, first observe that every \mathcal{Q} -category $X = (X, a)$ gives rise to the \mathcal{Q}^{op} -category $X^{\text{op}} = (X, a^\circ)$, where $a^\circ(x, y) = a(y, x)$

$(x, y \in X)$. With the commutative diagram

$$\begin{array}{ccc}
 (\mathcal{Q}\text{-}\mathbf{Chu})^{\text{op}} & \xrightarrow{(-)^{\text{op}}} & \mathcal{Q}^{\text{op}}\text{-}\mathbf{Chu} \\
 \text{cod}^{\text{op}} \downarrow & & \downarrow \text{dom} \\
 \mathcal{Q}\text{-}\mathbf{Cat} & \xrightarrow{(-)^{\text{op}}} & \mathcal{Q}^{\text{op}}\text{-}\mathbf{Cat}
 \end{array}$$

one sees that, up to functorial isomorphisms, $\text{cod}^{\text{op}} : (\mathcal{Q}\text{-}\mathbf{Chu})^{\text{op}} \rightarrow \mathcal{Q}\text{-}\mathbf{Cat}$ coincides with the small-topological functor $\text{dom} : \mathcal{Q}^{\text{op}}\text{-}\mathbf{Chu} \rightarrow \mathcal{Q}^{\text{op}}\text{-}\mathbf{Cat}$. \square

Corollary 3.6. $\mathcal{Q}\text{-}\mathbf{Chu}$ is complete and cocomplete, all small limits and colimits in $\mathcal{Q}\text{-}\mathbf{Chu}$ are preserved by both dom and cod .

Proof. The dom -initial lifting of a dom -structured limit cone in $\mathcal{Q}\text{-}\mathbf{Cat}$ is a limit cone in $\mathcal{Q}\text{-}\mathbf{Chu}$, which is trivially preserved. Having a right adjoint, dom also preserves all colimits. \square

Remark 3.7. (1) Let us describe (small) products in $\mathcal{Q}\text{-}\mathbf{Chu}$ explicitly: Given a family of \mathcal{Q} -distributors $\varphi_i : X_i \multimap Y_i$ ($i \in I$), one first forms the product X of the \mathcal{Q} -categories $X_i = (X_i, a_i)$ as in Remark 2.4(1) with projections p_i and the coproduct of the $Y_i = (Y_i, b_i)$ as in Remark 2.4(2) with injections s_i ($i \in I$). The transposes $\tilde{\varphi}_i$ then determines a \mathcal{Q} -functor $\tilde{\varphi}$ making the left square of

$$\begin{array}{ccc}
 \mathbf{P}X & \xleftarrow{p_i^*} & \mathbf{P}X_i \\
 \tilde{\varphi} \uparrow & & \uparrow \tilde{\varphi}_i \\
 Y & \xleftarrow{s_i} & Y_i
 \end{array}
 \qquad
 \begin{array}{ccc}
 X & \xrightarrow{(p_i)_\natural} & X_i \\
 \varphi \downarrow & & \downarrow \varphi_i \\
 Y & \xrightarrow{(s_i)_\natural} & Y_i
 \end{array}$$

commutative, while the right square exhibits φ as a product of $(\varphi_i)_{i \in I}$ in $\mathcal{Q}\text{-}\mathbf{Chu}$ with projections (p_i, s_i) , ($i \in I$); explicitly,

$$\varphi(x, y) = (\tilde{\varphi}(y))_x = (\tilde{\varphi}_i(y) \circ (p_i)_\natural)_x = (\tilde{\varphi}_i(y))_{p_i(x)} = \varphi_i(x_i, y)$$

for $x = ((x_i)_{i \in I}, q)$ in X and $y = s_i(y)$ in Y_i , $i \in I$.

- (2) The coproduct of $\varphi_i : X_i \multimap Y_i$ ($i \in I$) in $\mathcal{Q}\text{-}\mathbf{Chu}$ is formed like the product, expect that the roles of domain and codomain need to be interchanged. Hence, one forms the coproduct X of $(X_i)_{i \in I}$ and the product Y of $(Y_i)_{i \in I}$ in $\mathcal{Q}\text{-}\mathbf{Cat}$ and obtains the coproduct $\varphi : X \multimap Y$ in $\mathcal{Q}\text{-}\mathbf{Chu}$ as in

$$\begin{array}{ccc}
 X_i & \xrightarrow{(s_i)_\natural} & X \\
 \varphi_i \downarrow & & \downarrow \varphi \\
 Y_i & \xrightarrow{(\varphi_i)_\natural} & Y
 \end{array}$$

so that $\varphi(x, y) = \varphi_i(x, y_i)$ for $y = ((y_i)_{i \in I}, q)$ in Y and $x = s_i(x)$ in X_i , $i \in I$.

- The equalizer of $(f, g), (\bar{f}, \bar{g}) : \varphi \rightarrow \psi$ in $\mathcal{Q}\text{-}\mathbf{Chu}$ is obtained by forming the equalizer and coequalizer

$$U \xrightarrow{i} X \begin{array}{c} \xrightarrow{f} \\ \xrightarrow{\bar{f}} \end{array} Y \quad \text{and} \quad W \begin{array}{c} \xrightarrow{g} \\ \xrightarrow{\bar{g}} \end{array} Z \xrightarrow{p} V$$

in $\mathcal{Q}\text{-Cat}$, respectively. With $\tilde{\chi} : V \rightarrow \mathbf{P}U$ obtained from the coequalizer property making

$$\begin{array}{ccc} \mathbf{P}U & \xleftarrow{i^*} & \mathbf{P}X \\ \tilde{\chi} \uparrow & & \uparrow \tilde{\varphi} \\ V & \xleftarrow{p} & W \end{array}$$

commutative, Theorem 3.4 guarantees that

$$\chi \xrightarrow{(i,p)} \varphi \begin{array}{c} \xrightarrow{(f,g)} \\ \xrightarrow{(\bar{f},\bar{g})} \end{array} \psi$$

is an equalizer diagram in $\mathcal{Q}\text{-Chu}$, where

$$\chi(x, p(w)) = (\tilde{\chi}(p(w)))_x = (i^*(\tilde{\varphi}(w)))_x = (\tilde{\varphi}(w) \circ i_{\natural})_x = \varphi(i(x), w)$$

for all $x \in U$, $w \in W$.

- Coequalizers in $\mathcal{Q}\text{-Chu}$ are formed like equalizers, except that the roles of domain and codomain need to be interchanged.

We will now strengthen Corollary 3.6 and show total completeness and total cocompleteness of $\mathcal{Q}\text{-Chu}$ with the help of Proposition 2.6. To that end, let us observe that, since the limit and colimit preserving functors dom and cod must in particular preserve both monomorphisms and epimorphisms, a monomorphism $(f, g) : \varphi \rightarrow \psi$ in $\mathcal{Q}\text{-Chu}$ must be given by a monomorphism f and an epimorphism g in $\mathcal{Q}\text{-Cat}$, i.e., by an injective \mathcal{Q} -functor f and a surjective \mathcal{Q} -functor g . Consequently, $\mathcal{Q}\text{-Chu}$ is wellpowered, and so is its dual $(\mathcal{Q}\text{-Chu})^{\text{op}} \cong \mathcal{Q}^{\text{op}}\text{-Chu}$. The main point is therefore for us to prove:

Theorem 3.8. *$\mathcal{Q}\text{-Chu}$ contains a generating set of objects and, consequently, also a cogenerating set.*

Proof. With the notations explained below, we show that

$$\{\eta_s : \emptyset \twoheadrightarrow D_s \mid s \in \text{ob } \mathcal{Q}\} \cup \{\lambda_t : \{t\} \twoheadrightarrow \widehat{C} \mid t \in \text{ob } \mathcal{Q}\}$$

is generating in $\mathcal{Q}\text{-Chu}$. Here D_s belongs to a generating set of $\mathcal{Q}\text{-Cat}$ (see Remark 2.8), and

$$C = \coprod_{t \in \text{ob } \mathcal{Q}} \mathbf{P}\{t\}$$

is a coproduct in $\mathcal{Q}\text{-Cat}$ (see Remark 2.4(2)) of the presheaf \mathcal{Q} -categories of the singleton \mathcal{Q} -categories $\{t\}$ (see Proposition 3.2). From C one obtains \widehat{C} by adding an isomorphic copy of each object in C , which may be easily explained for a \mathcal{Q} -category (X, a) : simply provide the set $\widehat{X} := X \times \{1, 2\}$ with the structure

$$|(x, i)|_{\widehat{X}} = |x|_X \quad \text{and} \quad \widehat{a}((x, i), (y, j)) = a(x, y)$$

for all $x, y \in X$, $i, j \in \{1, 2\}$. Noting that the objects of $\mathbf{P}\{t\}$ are simply \mathcal{Q} -arrows with domain t , we now define $\lambda_t : \{t\} \twoheadrightarrow \widehat{C}$ by

$$\lambda_t(u, i) = \begin{cases} u & \text{if } \text{dom } u = t, \\ \perp & \text{else} \end{cases}$$

for $i \in \{1, 2\}$ and every object t and arrow u in \mathcal{Q} . For another element (v, j) in \widehat{C} , if $\text{dom } v = \text{dom } u = t$ one then has

$$[(u, i), (v, j)] \circ \lambda_t(u, i) = (v \swarrow u) \circ u \leq v = \lambda_t(v, j),$$

and in other cases this inequality holds trivially. Hence, λ_t is indeed a \mathcal{Q} -distributor.

Let us now consider \mathcal{Q} -Chu transforms $(f, g) \neq (\bar{f}, \bar{g}) : \varphi \rightarrow \psi$ as in

$$\begin{array}{ccc} (X, a) & \begin{array}{c} \xrightarrow{f^{\natural}} \\ \xrightarrow{\bar{f}^{\natural}} \end{array} & (Y, b) \\ \downarrow \varphi \circ & & \downarrow \psi \circ \\ (W, d) & \begin{array}{c} \xrightarrow{g^{\natural}} \\ \xrightarrow{\bar{g}^{\natural}} \end{array} & (Z, c) \end{array}$$

Case 1: $X = \emptyset$ is the initial object of $\mathcal{Q}\text{-Cat}$ (and $\mathcal{Q}\text{-Dis}$). Then $g \neq \bar{g}$, and we find $s \in \text{ob } \mathcal{Q}$ and $h : W \rightarrow D_s$ with $hg \neq h\bar{g}$ in $\mathcal{Q}\text{-Cat}$. Consequently, $(1_{\emptyset}, h) : \eta_s \rightarrow \varphi$ satisfies $(f, g)(1_{\emptyset}, h) \neq (\bar{f}, \bar{g})(1_{\emptyset}, h)$.

Case 2: $f \neq \bar{f}$, so that $f(x_0) \neq \bar{f}(x_0)$ for some $x_0 \in X$. Then, for $t := |x_0|$, $e : \{t\} \rightarrow X$, $|x_0| \mapsto x_0$, is a \mathcal{Q} -functor with $fe \neq \bar{f}e$, and it suffices to show that

$$h : W \rightarrow \widehat{C}, \quad w \mapsto (\varphi(x_0, w), 1)$$

is a \mathcal{Q} -functor making $(e, h) : \lambda_t \rightarrow \varphi$ a \mathcal{Q} -Chu transform. Indeed,

$$\begin{aligned} d(w, w') &\leq \varphi(x_0, w') \swarrow \varphi(x_0, w) = [h(w), h(w')], \\ \lambda_t(h(w)) &= \varphi(x_0, w) = \varphi(e(t), w) \end{aligned}$$

for all $w, w' \in W$.

Case 3: $X \neq \emptyset$ and $g \neq \bar{g}$. Then $g(z_0) \neq \bar{g}(z_0)$ for some $z_0 \in Z$, and with any fixed $x_0 \in X$ we may alter the previous definition of $h : W \rightarrow \widehat{C}$ by

$$h(w) := \begin{cases} (\varphi(x_0, w), 2) & \text{if } w = \bar{g}(z_0), \\ (\varphi(x_0, w), 1) & \text{else.} \end{cases}$$

The verification for h to be a \mathcal{Q} -functor and $(e, h) : \lambda_t \rightarrow \varphi$ a \mathcal{Q} -Chu transform remain intact, and since $hg \neq h\bar{g}$, the proof is complete. \square

Remark 3.9. A generating set in $\mathcal{Q}\text{-Chu}$ may be alternatively given by

$$\{\lambda_{\emptyset} : \emptyset \rightarrow \widehat{C}\} \cup \{\lambda_t : \{t\} \rightarrow \widehat{C} \mid t \in \text{ob } \mathcal{Q}\},$$

so that in Case 1 one may proceed exactly as in Case 3 only by replacing $\varphi(x_0, w)$ with $\top : q \rightarrow |w|$ for any fixed $q \in \text{ob } \mathcal{Q}$.

With Theorem 3.8 we obtain:

Corollary 3.10. *$\mathcal{Q}\text{-Chu}$ is totally complete and totally cocomplete and, in particular, hypercomplete and hypercomplete.*

REFERENCES

- [1] J. Adámek. Colimits of algebras revisited. *Bulletin of the Australian Mathematical Society*, 17:433–450, 1977.
- [2] J. Adámek, H. Herrlich, and G. E. Strecker. *Abstract and Concrete Categories: The Joy of Cats*. Wiley, New York, 1990.
- [3] M. Barr. *-Autonomous categories and linear logic. *Mathematical Structures in Computer Science*, 1:159–178, 1991.
- [4] J. Barwise and J. Seligman. *Information Flow: The Logic of Distributed Systems*, volume 44 of *Cambridge Tracts in Theoretical Computer Science*. Cambridge University Press, Cambridge, 1997.
- [5] J. Bénabou. Les distributeurs. Université Catholique de Louvain, Institute de Matématique Pure et Appliquée, Rapport no. 33, 1973.
- [6] J. Bénabou. Distributors at work. Lecture notes of a course given at TU Darmstadt, 2000.
- [7] F. Borceux. *Handbook of Categorical Algebra: Volume 1, Basic Category Theory*, volume 50 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1994.
- [8] R. Börger and W. Tholen. Cantors Diagonalprinzip für Kategorien. *Mathematische Zeitschrift*, 160(2):135–138, 1978.
- [9] R. Börger and W. Tholen. Total categories and solid functors. *Canadian Journal of Mathematics*, 42:213–229, 1990.

- [10] B. Ganter. Relational Galois connections. In S. O. Kuznetsov and S. Schmidt, editors, *Formal Concept Analysis*, volume 4390 of *Lecture Notes in Computer Science*, pages 1–17. Springer, Berlin-Heidelberg, 2007.
- [11] E. Giuli and W. Tholen. A topologist’s view of Chu spaces. *Applied Categorical Structures*, 15(5-6):573–598, 2007.
- [12] M. Grandis. Weak subobjects and the epi-monic completion of a category. *Journal of Pure and Applied Algebra*, 154(1-3):193–212, 2000.
- [13] M. Grandis. On the monad of proper factorisation systems in categories. *Journal of Pure and Applied Algebra*, 171(1):17–26, 2002.
- [14] H. Heymans. *Sheaves on Quantales as Generalized Metric Spaces*. PhD thesis, Universiteit Antwerpen, Belgium, 2010.
- [15] D. Hofmann, G. J. Seal, and W. Tholen, editors. *Monoidal Topology: A Categorical Approach to Order, Metric, and Topology*, volume 153 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 2014.
- [16] U. Höhle and T. Kubiak. A non-commutative and non-idempotent theory of quantale sets. *Fuzzy Sets and Systems*, 166:1–43, 2011.
- [17] G. M. Kelly. *Basic Concepts of Enriched Category Theory*, volume 64 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 1982.
- [18] F. W. Lawvere. Metric spaces, generalized logic and closed categories. *Rendiconti del Seminario Matematico e Fisico di Milano*, XLIII:135–166, 1973.
- [19] S. G. Matthews. Partial metric topology. *Annals of the New York Academy of Sciences*, 728(1):183–197, 1994.
- [20] V. Pratt. Chu spaces and their interpretation as concurrent objects. In J. Leeuwen, editor, *Computer Science Today*, volume 1000 of *Lecture Notes in Computer Science*, pages 392–405. Springer, Berlin-Heidelberg, 1995.
- [21] Q. Pu and D. Zhang. Preordered sets valued in a GL-monoid. *Fuzzy Sets and Systems*, 187(1):1–32, 2012.
- [22] K. I. Rosenthal. *The Theory of Quantaloids*, volume 348 of *Pitman Research Notes in Mathematics Series*. Longman, Harlow, 1996.
- [23] L. Shen and W. Tholen. Topological categories, quantaloids and Isbell adjunctions. *arXiv:1501.00703*, 2015.
- [24] R. Street and R. F. C. Walters. Yoneda structures on 2-categories. *Journal of Algebra*, 50(2):350–379, 1978.
- [25] I. Stubbe. Categorical structures enriched in a quantaloid: categories, distributors and functors. *Theory and Applications of Categories*, 14(1):1–45, 2005.
- [26] I. Stubbe. Categorical structures enriched in a quantaloid: tensored and cotensored categories. *Theory and Applications of Categories*, 16(14):283–306, 2006.
- [27] I. Stubbe. An introduction to quantaloid-enriched categories. *Fuzzy Sets and Systems*, 256(0):95–116, 2014. Special Issue on Enriched Category Theory and Related Topics (Selected papers from the 33rd Linz Seminar on Fuzzy Set Theory, 2012).
- [28] Y. Tao, H. Lai, and D. Zhang. Quantale-valued preorders: Globalization and cocompleteness. *Fuzzy Sets and Systems*, 256(0):236–251, 2014. Special Issue on Enriched Category Theory and Related Topics (Selected papers from the 33rd Linz Seminar on Fuzzy Set Theory, 2012).
- [29] W. Tholen. Semi-topological functors I. *Journal of Pure and Applied Algebra*, 15(1):53–73, 1979.
- [30] W. Tholen. Note on total categories. *Bulletin of the Australian Mathematical Society*, 21:169–173, 1980.
- [31] R. F. C. Walters. Sheaves and Cauchy-complete categories. *Cahiers de Topologie et Géométrie Différentielle Catégoriques*, 22(3):283–286, 1981.

LILI SHEN, DEPARTMENT OF MATHEMATICS AND STATISTICS, YORK UNIVERSITY, TORONTO, ONTARIO, CANADA, M3J 1P3

E-mail address: shenlili@yorku.ca

WALTER THOLEN, DEPARTMENT OF MATHEMATICS AND STATISTICS, YORK UNIVERSITY, TORONTO, ONTARIO, CANADA, M3J 1P3

E-mail address: tholen@mathstat.yorku.ca